

## Response of *Physalis peruviana* L. genotypes to *Fusarium oxysporum* f. sp. *physali* under greenhouse

### Respuesta de genotipos de *Physalis peruviana* L. A *Fusarium oxysporum* f. sp. *physali* bajo invernadero

Johana Pabón-Villalobos<sup>1</sup>; Tulio César Lagos-Burbano<sup>2</sup>; Diego Fernando Mejía-España<sup>3</sup>; David Duarte-Alvarado<sup>4</sup>

#### ARTICLE DATA

<sup>1</sup> Researcher, M.Sc. Grupo de Investigación en Producción de Frutales Andinos (GPFA), Universidad de Nariño, Pasto, Colombia, [johanapv1125@udenar.edu.co](mailto:johanapv1125@udenar.edu.co)  
<https://orcid.org/0000-0002-0028-5603>

<sup>2</sup> Professor, PhD. Grupo de Investigación en Producción de Frutales Andinos (GPFA), Universidad de Nariño, Pasto, Colombia, [tclagosb@udenar.edu.co](mailto:tclagosb@udenar.edu.co)  
<https://orcid.org/0000-0001-9222-4674>

<sup>3</sup> M.Sc. Grupo de Investigación Tecnologías Emergentes en Agroindustria (TEA), Universidad de Nariño, Pasto, Colombia, [diegomejias@udenar.edu.co](mailto:diegomejias@udenar.edu.co)  
<https://orcid.org/0000-0002-6707-5803>

<sup>4</sup> M.Sc. Grupo de Investigación en Producción de Frutales Andinos (GPFA), Universidad de Nariño, Pasto, Colombia, [david890223@udenar.edu.co](mailto:david890223@udenar.edu.co)  
<https://orcid.org/0000-0002-8896-3594>

**Cite:** Pabón-Villalobos, J.; Lagos-Burbano, T.; Mejía-España, D.; Duarte-Alvarado, D. (2022). Response of *Physalis peruviana* L. genotypes to *Fusarium oxysporum* f. sp. *physali* under greenhouse. *Revista de Ciencias Agrícolas*. 39(2): 89-107.  
doi: <https://doi.org/10.22267/rcia.223902.185>

Received: October 06 2022.

Accepted: September 15 2022.



#### ABSTRACT

The goldenberry (*Physalis peruviana*) is an exotic fruit that in recent years has acquired great importance in both the local and international markets; one of the limiting phytosanitary problems for this crop is vascular wilt caused by *Fusarium oxysporum* f. sp. *physali*, which causes losses of 80-90%. The management of this pathogen is difficult and so far, it is based on preventive measures; however, there are alternatives such as genetic resistance, which is one of the most effective and profitable measures for its management. Taking that into account, the objective of this study was to evaluate the reaction of 40 genotypes of goldenberry against *F. oxysporum* under greenhouse conditions, by means of pathogenicity tests. The experiment was conducted in a selected place in the city of Pasto (Nariño department, south of Colombia). It was carried out with 40 genetic materials corresponding to different genotypes, one commercial control and four replicates per experimental unit; the statistical design was completely randomized. The traits evaluated were plant height (cm), disease severity (%), AUDPC area under the disease progress curve (units), disease incidence (%) and degree of vascular discoloration. The genotypes 09U138 and 12U399 have greater plant height (50.19 and 47.36 cm), lower AUDPC (zero units), lower incidence (0%) and lower degree of vascular discoloration (zero), with statistical differences from the rest of the genotypes, including the control. Field evaluations should be conducted with the same isolation and other commercial controls, as this research is only a step forward in the search for the resistance of uchuva to *F. oxysporum*.

**Keywords:** Uchuva; Goldenberry; losses; pathogenicity; resistance; vascular wilt.

#### RESUMEN

La uchuva (*Physalis peruviana*) es una fruta exótica que ha adquirido gran importancia tanto en el mercado local como internacional. Uno de los problemas fitosanitarios limitantes para este cultivo es la marchitez vascular causada por *Fusarium oxysporum* f. sp. *physali*, que provoca

pérdidas del 80-90%. El manejo de este patógeno es difícil y hasta el momento, se basa en medidas preventivas; sin embargo, existen alternativas como la resistencia genética, que es una de las medidas más efectivas y rentables para su manejo. El objetivo de este estudio fue evaluar mediante pruebas de patogenicidad y bajo condiciones de invernadero, la reacción de 40 genotipos de uchuva frente a una cepa de *Fusarium oxysporum*. El experimento que se realizó en Colombia en una localidad de Pasto (Nariño) constó de 39 materiales genéticos, un testigo comercial y cuatro repeticiones por unidad experimental. Se utilizó el Diseño Completamente al Azar. Las características evaluadas fueron altura de planta (cm), severidad (%), área bajo la curva de progreso de la enfermedad (ABCPE), incidencia de la enfermedad (%) y el grado de decoloración vascular. Los genotipos 09U138 y 12U399 presentaron mayor altura de planta (50,19 y 47,36cm), menor ABCPE, menor incidencia y menor grado de decoloración vascular, con diferencias estadísticas con respecto al resto de los genotipos, incluido el testigo. Se deben realizar evaluaciones de campo con el mismo aislamiento y otros testigos comerciales, pues esta investigación es solo un avance en la búsqueda de la resistencia de uchuva a *F. oxysporum*.

**Palabras clave:** uchuva; Goldenberry; pérdidas; patogenicidad; resistencia; marchitez vascular.

## INTRODUCTION

The goldenberry, or also known as Goldenberry or Uchuva (*Physalis peruviana* L.), is an Andean fruit species that has become an alternative for the economy of many countries as it stands out as an export product; its importance derives from the nutritional characteristics and medicinal properties (Chávez *et al.*, 2019). In Colombia, cape gooseberry is the second most exported product after bananas; its cultivation offers great advantages because, being Colombia a tropical country, the permanent production of the fruit can be guaranteed for international markets (Ruiz *et al.*, 2018; García *et al.*, 2021). In Colombia, the producing departments are Boyacá, Cundinamarca, Antioquia, Nariño, Norte de Santander, Santander, Huila, Tolima, Cauca and Valle del Cauca, with Boyacá being the largest producer with 30.6% of the planted area, followed by Cundinamarca with 28.6% and Nariño with 13.4% (EVA, 2021).

Among the most important diseases of the crop, we can find vascular wilt (*Fusarium oxysporum* f. sp. *physali*), dieback (*Phoma physalidis*), gray leaf spot (*Cercospora physalidis*), fruit spot (*Alternaria* sp.), and Gray mold (*Botrytis cinerea*); other diseases reported in Colombia, but of lower incidence

are cottony rot (*Sclerotinia sclerotiorum*), stem rot (*Pythium* sp.), Anthracnose (*Colletotrichum gloeosporioides*), the bacteria *Xanthomonas* sp., *Ralstonia solanacearum*, nematodes such as *Meloidogyne* sp., *Pratylenchus* sp., and viruses such as leaf-roll virus (PLRV), Potato virus Y (PVY) and Andean mottle virus (APMV) (Ruiz *et al.*, 2018; Diaz *et al.*, 2019).

Vascular wilt caused by *Fusarium oxysporum* f. sp. *physalis* (*FoPh*) is one of the most limiting diseases for Goldenberry as it has generated losses between 80 and 90% (García *et al.*, 2021; Simbaqueda *et al.*, 2021). In the last decade, the department of Cundinamarca suffered at great loss of the crop and it was necessary to relocate it to Boyacá department; that is why the production of this fruit is concentrated in this place (Valderrama, 2018; Simbaqueba *et al.*, 2018; Chávez *et al.*, 2019; Simbaqueba *et al.*, 2021). This fungus is difficult to manage due to soil contamination with the pathogen, something that occurs when harvest residues and affected plant tissues are not properly discarded, or when contaminated soil is moved (Valderrama, 2018). Plants can become infected with the fungal reproductive structures such as mycelium, conidia, and chlamydospores (resistance structures that manage to remain up to thirty years in soils), which germinate

upon contact with host plant root exudates (Vásquez & Castaño, 2017; Joshi, 2018; Giraldo *et al.*, 2020). Plants affected by FoPh are initially characterized by leaf chlorosis followed by generalized yellowing. Symptoms that are mixed with loss of turgor in branches and stems. In general, these symptoms tend to be unilateral, affecting one or two of the main stems; it is also common to see that they are bent leaving the fruits attached to them.

The progression of the disease ends up affecting the entire plant which eventually dies (Gordon, 2017; Gonzáles, 2019; Chávez *et al.*, 2019; Giraldo *et al.*, 2020) When longitudinal cuts are made in the stems and branches, a light brown coloration of the parenchyma is observed, which in advanced stages turns brown; comparatively, the bark tissues are seemingly healthy (Valderrama, 2018; Agudelo, 2020).

Disease management is based on preventive measures that include not planting in plots with a history of incidence by *F. oxysporum*, the use of pathogen-free propagation material, avoiding unnecessary wounds during cultivation, weed control to reduce excess moisture, avoiding soil waterlogging, eradication of diseased plants, crop rotation, and solarization (Moreno *et al.*, 2019; Berruezo, 2018; Chávez *et al.*, 2020). Non-preventive alternatives include biological control and the search for resistant cultivars; the latter is seen as one of the most effective and economically profitable measures for disease management in the field (Vásquez & Castaño, 2017; Rodríguez & Pedraza, 2019).

Regarding the resistance response to Foph to avoid losses in the producing areas, numerous efforts have been made, such as the one done by Pulido *et al.* (2011), who carried out bioassays with 70 goldenberry accessions

and through cluster analysis, concluded that three of the evaluated materials presented resistance to the pathogen. Rodríguez (2013), on the other hand, through pathogenicity tests in the field, identified two introductions of *Physalis* with significant values of resistance to vascular wilt. Osorio *et al.* (2016) identified promising accessions with different degrees of resistance, as well as 16 markers associated with the resistance response. Mayorga *et al.* (2019) found genetic materials with desirable agronomic traits and excellent response to Foph attack and considered them important for breeding schemes.

Until 2018, the department of Nariño did not present a historical record of incidence of vascular wilt (Agencia UNAL, 2018). However, foci of the disease have been identified, so the efforts of producers and technicians should be oriented to avoid its appearance and dissemination, due to the considerable losses in yields that this fungus causes.

The objective of this work was to evaluate the reaction of Goldenberry genotypes against *Fusarium oxysporum* f. sp. *physali* (FoPh) under greenhouse conditions, with the purpose of identifying sources of resistance that could be used in breeding programs for the production of improved cultivars with resistance to the disease.

## MATERIALS AND METHODS

**Location.** The evaluation of the 40-goldenberry genotypes and their reaction to *Fusarium oxysporum* attack was carried out in the greenhouse located in the Agrosavia C.I. Obonuco facilities, at 2760masl and with an average temperature of 13°C.

**Genetic materials.** Forty genetic materials were used; 9 double haploid lines and 11 genotypes of the *Fusarium* group belong to the Colombian Agricultural Research Corporation AGROSAVIA Tibaitatá, 19 genotypes from the University of Nariño and a commercial control, which is a selection made by cape gooseberry farmers in the department of Nariño (Table 1). The seeds were initially sown in germination trays with peat substrate. When the seedlings presented 3 to 4 true leaves, they were taken to 1kg bags containing sterile soil and irrigated four days a week for 4 hours by mist irrigation. Fertilization was edaphic.

During transplantation and growth, a mixture of DAP+ Agriminis® (0.5g/plant) was applied, during the flowering stage a mixture of calcium nitrate+10-30-10 (2g/plant) was used and during production, potassium nitrate (3g/plant). Monthly applications of 15-15-15 (5g/plant) were then made, which were increased over time. A V-shaped trellis system was used; with pruning every 20 days, weed control was manual and harvests were scheduled twice a month. To control the stem borer, Exalt® was applied at the dose suggested by the manufacturer (2mL/L).

### Isolation and purification of the pathogen.

The pathogen strain used in this study was obtained from one of the experimental plots located in Puerres. For its isolation, samples were taken from plant stems showing vascular necrosis; diseased tissue cuts of approximately 3 mm were made, and each cut underwent the disinfection protocol (1% sodium hypochlorite, 70% alcohol, and sterile distilled water washings). The disinfected tissues were seeded in PDA culture medium (39g/L of water) and incubated for eight days at 22°C. Colonies that presented a whitish cottony appearance and pink color on the underside were replicated in the same medium for isolation and subsequent identification. A small amount of mycelium was then extracted with the aid of a dissecting needle and observed under the microscope at 40X, identifying microconidia with 0 to 3 septate macroconidia characteristic of *Fusarium oxysporum* (Carmona *et al.*, 2020). The isolation used was molecularly confirmed by DNA sequencing of the ITS and EF1 $\alpha$  genes carried out in the molecular biology laboratories of Agrosavia at C.I. Tibaitatá.

**Table 1.** Genetic materials evaluated for their reaction against to *Fusarium oxysporum* f.sp. *physali*.

AGROSAVIA		UDENAR	
12U347	09U086	UN01	NEIRA
12U350	09U089	UN03	UN45
12U352	09U099	UN13	UN49
12U357	09U116	UN14	UN52
12U360	09U128	UN19	SILVANIA
12U368	09U136	UN26	PERU
12U374	09U138	UN30	COLOMBIA
12U377	09U140	UN34	PURACÉ
12U399	13U407	UN35	KENIA
ANDINA	13U408	UN36	CONTROL

**Inoculation by root immersion.** The inoculum was obtained in PDA culture medium; the Petri dishes seeded with the pathogen were incubated for eight days. The conidia were removed with the help of a bacteriological rake by adding 200mL of sterile distilled water with 0.1% Tween 80, then filtered through a gauze. The suspension was adjusted to  $1 \times 10^6$  conidia/mL calibrated using a hemacytometer, following the methodology described by Arellano (2018) and Agudelo (2020). When the goldenberry plants were 20 cm high, they were removed from their plastic bags; the roots were washed with sterile distilled water and with the help of scissors disinfested with 1%, quaternary ammonium. The apexes of the main root were cut; then they were submerged from the stalk base to the root for 30 minutes in 250 mL plastic cups containing the conidial suspension. Finally, the plants were planted in bags with sterilized soil. Arellano (2018), Ángel *et al.* (2018) and Agudelo (2020) have used this method.

## Traits evaluated

**Plant height (cm).** The height of each genotype and its replicates were recorded weekly by using a tape measure.

**Severity (%).** Severity (Table 2) was recorded every eight days, using the scale proposed by Garcés *et al.* (2017).

**Area under the disease progress curve (AUDPC).** To compare the differences between genetic materials from disease severity values, the AUDPC calculation was performed, this parameter incorporated the speed of disease progression and severity into a single value. That is, accumulation of daily values of the percentage of infection interpreted directly without performing any transformation (Chañag *et al.*, 2017; Sánchez *et al.*, 2017; Bocianowski *et al.*, 2020).

**Table 2.** Scale used to qualify the degree of severity of vascular wilt caused by *Fusarium oxysporum*.

Severity	Percentage	Traits
0	0	No symptoms manifested
1	10	Mild chlorosis, without necrotic lesions or more than 10% of the total foliage is wilted and/or chlorotic.
3	25	Leaves wilted and/or chlorotic
5	50	50% of the leaves show the beginnings of wilting
7	75	75% of foliage severely wilted
9	100	Dead or severely infected plants showing practically all their foliage wilted, with chlorosis, necrosis, and/or premature defoliation.

$$AUDPC = \sum_{t=1}^n \frac{(X_{i+1} + X_i)}{2} * (T_{i+1} - T_i)$$

Where:  $X_i$  = Proportion of affected tissue under observation  $i$ ,  $T_{i+1}-T_i$  = time in days between two readings,  $n$  = Total number of observations.

**Incidence (%).** The number of diseased plants was evaluated weekly- The incidence was obtained by applying the formula:

$$\text{Incidence (\%)} = [\text{Number of sick individuals} / \text{total individuals}] \times 100$$

**Vascular discoloration.** Once the observations were finished, the plants were removed from the plastic bags and a transversal cut was made; the upper, middle, and lower part of each genotype was evaluated, using the scale (Table 3) proposed by Garcés *et al.* (2017).

**Susceptibility index.** The mean and standard deviation statistics were used; then a relative value was assigned to the evaluated traits to obtain a score that allowed a classification of the tested genotypes.





**Experimental design and statistical analysis.** The Completely Randomized Design

was used with 39 genotypes, a commercial control and five plants like replications per experimental unit, of which one was a control inoculated with water. During the entire experiment, 28 evaluations were made; it should be noted that after reading 14, the surviving genotypes were inoculated again to verify that the resistance presented was not associated with escape. Analysis of Variance (ANOVA) and Tukey's mean comparison tests at 95% probability were carried out using the statistical program Statgraphics Centurion XVII.

## RESULTS AND DISCUSSION

**Plant height.** The ANOVA for plant height indicated significant statistical differences among the genetic materials evaluated ( $P < 0.05$ ). The Tukey test for comparison of means showed that the genotype that presented the greatest growth was 12U399 with 49.94cm, followed by 09U138 with 46.73cm; results that did not differ statistically from the commercial control (35.09cm), but from UN35 that showed the lowest average (9.61cm); 13U407 and UN34 showed heights of 44.73, 44.21, 43.92 and 41.35cm respectively, with no statistical difference with each other; the other plants showed averages fluctuating between 40.91 and 22.64cm.

**Table 3.** Vascular discoloration scale for vascular wilt caused by *F. oxysporum*.

Scale	Description	Color
0	None	
1	Slight	
2	Intermediate	
3	Severe	

On the other, the growth for 09U136, 12U374, Puracé, and UN49 was 18.67, 18.22, 17.15, and 14.32cm. In that regard, Insuasty *et al.* (2014) obtained low height averages by studying the pathosystem pea-*Fusarium oxysporum*, and they concluded that growth was reduced due to the pathogen's ability to colonize roots, which in turn prevents adequate nutrition; Wang & Jeffers (2002) state that hostas (*Hosta* spp.) plants, when inoculated with strains of *Fusarium otae* experienced growth retardation, Rose *et al.* (2003) reported that squash (*Cucurbita pepo*) plants infected with *Fusarium oxysporum* f.sp *radiciscucumerinum* showed a lower height compared to the commercial control, thus showing the negative effect of the pathogen on this trait.

Alvarado (2005), when evaluating pathogenic strains of *Fusarium* sp. in colored calla lilies (*Zantedeschia* spp.) indicates that one of the most important aspects affecting this fungus is size; at the end of the experiment, the calla lilies, besides being dwarf, had small and thin leaves, showing poor root development. *F. oxysporum* is a fungus that invades the cortical cells of the root intercellularly during the infection process; it enters the vascular system through the xylem and begins to produce microconidia, which rapidly colonize the plant generating obstructions by hyphae, which prevent communication between the cells of the conducting vessels, blocking the flow of water and nutrients and directly interfering with plant growth (Marín *et al.*, 2018; Chávez *et al.*, 2019; Srinivas *et al.*, 2019; Giraldo *et al.*, 2020).

The results obtained for this trait (Table 4) in genotypes 12U399 (49.94cm) and 09U138 (46.73cm) indicate resistance to *FoPh*. In

this regard, Rodríguez (2020) states that in resistant crops, when the pathogen attack is detected, flavonoids such as catechins and their oxidation products inactivate the enzymes, so that colonization is confined to the initial infection points. Bani *et al.* (2018) point out that resistant plants are able to prevent the advance of the fungus by sealing the xylem by means of gels or gums constituted by polysaccharides of high molecular weight.

**Area under the disease progression curve (AUDPC).** From the severity data, the area under the disease progress curve (AUDPC) was obtained; value that in turn was calculated using the trapezoidal integration model (Alves *et al.*, 2017; Castellanos *et al.*, 2021), finding significant statistical differences ( $P < 0.05$ ) in disease intensity over time for the genotypes evaluated. Tukey's mean comparison test indicated that genotypes 09U138 and 12U399 differed from the others, even from the commercial control (975 units) by presenting the lowest AUDPC values with zero units (Table 4), followed by Peru with 529 units. UN34, 13U407, and 09U089 revealed AUDPC of 803, 805, and 815 respectively without statistical differences with each other; the rest of the genotypes showed AUDPC ranging from 828 to 1340 units.

The highest values (Table 4) were for 09U128, 09U086, 09U136, Puracé, UN49, and UN35 with 1398, 1400, 1403, 1428, 1473, and 1519 units, which were not different from the commercial control (975 units). Materials 09U138 and 12U399 were the least affected by the disease and apparently could be classified as resistant to *F. oxysporum* f. sp. *physali* attack.

It is worth mentioning that at the end of evaluation 14, a second inoculation of the surviving genotypes was performed by the root immersion method, as explained in the methodology, to verify that the absence of symptoms was not associated with escape.

The results validate the report by Osorio *et al.* (2017), who through conglomerate analysis identified the resistance of different Goldenberry accessions to FoPh attack, the genotype 09U138 was characterized by low severity and lower AUDPC.

**Table 4.** Tukey's means test for plant height, severity and incidence of the Goldenberry genotypes evaluated.

GENOTYPES	HEIGHT	GENOTYPES	AUDPC	GENOTYPES	INCIDENCE
12U399	49.94	A	12U399	0	A
09U138	46.73	AB	09U138	0	A
12U357	44.73	ABC	PERÚ	529	AB
NEIRA	44.21	ABC	UN34	803	ABC
13U407	43.92	ABC	13U407	805	ABC
UN34	41.35	ABC	09U089	815	ABC
12U377	40.91	ABCD	SILVANIA	828	BC
SILVANIA	40.56	ABCDE	UN26	880	BC
KENIA	38.82	ABCDEF	UN52	936	BC
UN52	38.47	ABCDEF	CONTROL	975	BC
09U089	37.62	ABCDEF	12U352	1031	BC
12U352	37.15	ABCDEF	UN36	1053	BC
12U350	37.01	ABCDEF	NEIRA	1071	BC
UN36	36.66	ABCDEF	12U368	1085	BC
UN45	36.36	ABCDEF	UN14	1090	BC
UN13	36.32	ABCDEF	ANDINA	1099	BC
UN26	35.78	ABCDEF	UN13	1123	BC
PERÚ	35.75	ABCDEF	COLOMBIA	1134	BC
CONTROL	35.09	ABCDEF	09U099	1148	BC
12U360	33.45	ABCDEF	UN45	1151	BC
12U347	33.10	ABCDEF	UN30	1155	BC
UN03	33.08	ABCDEF	UN19	1157	BC
UN19	32.86	ABCDEF	13U408	1172	BC
09U099	31.09	ABCDEF	KENIA	1179	BC
12U368	29.75	ABCDEF	12U360	1248	BC
UN01	29.70	ABCDEF	UN01	1256	BC
UN30	29.47	ABCDEF	12U347	1291	BC
09U128	29.32	ABCDEF	09U116	1305	BC
COLOMBIA	29.24	ABCDEF	12U377	1307	BC
13U408	29.05	ABCDEF	09U140	1307	BC
UN14	26.29	BCDEF	12U374	1311	BC
09U086	25.22	BCDEF	UN03	1318	BC
09U140	23.54	CDEF	12U350	1333	BC
09U116	23.18	CDEF	12U357	1340	BC
ANDINA	22.64	CDEF	09U128	1398	C
09U136	18.67	DEF	09U086	1400	C
12U374	18.22	EFG	09U136	1403	C
PURACE	17.15	FGH	PURACE	1428	C
UN49	14.32	GH	UN49	1473	C
UN35	9.61	H	UN35	1519	C

Means with different letters indicate significant difference according to Tukey test ( $P \leq 0.05$ ).




Regarding resistance, Pulido (2010) and Gayosso *et al.* (2021) explain that once pathogens overcome mechanical barriers to infection, plant receptors initiate signaling pathways that drive the expression of defense response genes that depend on their ability to recognize harmful molecules, carry out signal transduction, and respond defensively to a potential invader. Burbano (2020) and Kaur *et al.* (2021) additionally state that when a resistance gene identifies an avirulence gene in the pathogen, a process of cell death is activated in the infected cell, completely stopping colonization; thus, granting resistance to the plant, which is what possibly occurred with genotypes 09U138 and 12U399. However, it is recommended that these genetic materials be evaluated against other strains of the fungus. Pournalibaba *et al.* (2017), Bani (2018), and Joshi (2018) explain that *Fusarium oxysporum* colonization is restricted in cultivars resistant to the region of initial entry of the pathogen, due to occlusion of the vessels by gels, callose, and tyloses deposition.

The high AUDPC values observed in the rest of the genotypes apparently, indicated low amounts of callose and tyloses that were degraded by the pectolytic enzymes of the pathogen, which interfered in the activation of the defense system against *Fusarium oxysporum*, showing a high susceptibility and a compatible interaction where the pathogen initiates the infection in which it can multiply and progress systemically, invading other tissues triggering the disease (Chañang *et al.*, 2017; Castro *et al.*, 2020; Islas, 2021; Kaur *et al.*, 2021).

For infection to be successfully achieved, the pathogen-plant interaction responds to a process where different sets of genes, must be mobilized that allow early host signaling, adhesion to the host surface, enzymatic breakdown of physical barriers, defense against plant antifungal compounds, and inactivation and subsequent death of host cells by secreted mycotoxins (Agrios, 2005).

The expression of symptoms in susceptible genotypes was observed after the second week of inoculation with the fungus, results that coincide with those reported by Pulido (2010) and Osorio *et al.* (2017), who affirms that in the case of Goldenberry the age of the plant does not influence the speed of infection and symptoms begin to be evident within the first two weeks after inoculation.

Figure 1 shows the initial and final severity of each of the genotypes evaluated, it can be clearly seen that almost all reached a final affectation of 100%, except for 13U407, UN26, Sylvania, commercial control, UN14, UN34, and 12U352, whose severities ranged between  $74 \pm 24.7\%$  and  $78 \pm 24.7\%$ ; on the other hand, Peru, UN52, and 09U089 showed damages of 50, 53, and  $54 \pm 24.7\%$ , respectively, and genotypes 09U138 and 12U399, showed the lowest result for this trait (0%), which is lower than that of the commercial control ( $76 \pm 24.7\%$ ), which undoubtedly indicates resistance to pathogen attack; it was also possible to appreciate that the materials UN36, Andina, UN45, 09U116, 12U347, and 12U368 initially did not show symptoms associated with *FoPh*, but at the end of the experiment they showed 100% damage which was the highest value.



**Figure 1.** Initial and final severity for the 40 cape gooseberry genotypes inoculated with *Fusarium oxysporum* f. sp. *physalis*.

In this regard, Cubedo (2008) and Marín *et al.* (2018), mention that the diagnosis of *F. oxysporum* cannot be made immediately since the fungus colonizes the vascular system before the expression of symptoms in the plant, that is, when the disease is detected, the infection is already in advanced stage.

**Incidence.** The analysis of variance revealed significant differences between the evaluated genotypes ( $P < 0.05$ ), and the Tukey test for


comparison of means between genotypes showed that 09U138 and 12U399 had the lowest average for this trait, which was 0, results that differed statistically from the commercial control (75%), followed by Peru and 09U089 with incidences of 25 and 50% respectively, on the other hand the genotypes UN34, 12U352, UN14, 12U360, 09U099, Sylvania, 13U407, the commercial control, and UN52 had an average of 75% diseased plants with no statistical differences with

each other, even when compared to the control (75%), the rest of the genotypes with the exception of UN26 (81.25%), showed the highest percentage which was 100% (Table 4). When these plants were visually analyzed wilting was observed on the leaves, mainly in the lower part extending to the upper part, causing defoliation and necrosis (Figure 2), therefore they are considered susceptible to attack by *F. oxysporum* f. sp. *physali*, and their use is not recommended in soils where there is high inoculum pressure

The results show that these genetic materials lack effective defense mechanisms against the strain used, but their response to other strains is unknown; it should be noted that there are no treatments to cure plants infected by FoPh; besides, this fungus has a wide range of hosts, so measures such as crop rotation are not effective. The use of biological agents does not achieve the desired

success because they can be affected by biotic and abiotic factors that make biological control in the field inconsistent.

In this research the fungus was able to cause the disease, associated symptoms, and even plant death; in commercial plots, the use of these varieties could cause an increase in inoculum and spread of the pathogen. In relation to the results obtained in genotypes 09U138 and 12U399, it can be inferred that they showed resistance to the attack of *FoPh* from the first to the last day of the evaluations carried out, since no symptoms associated with the disease such as wilting of basal leaves, loss of turgor, epinasty, chlorosis, prostration of the stalk, and the stem, were observed during the first or second inoculation, chlorosis, stem prostration or necrosis of the stem, or roots (Figure 2 D), this is how these two genotypes become an option for the management of the fungus in our region.



A. Plant wilt. B. Chlorotic leaves. C. Loss of turgor. D. Healthy plant.

**Figure 2.** Symptoms of vascular wilt in Goldenberry caused by *Fusarium oxysporum* f. sp. *physalis*.


Controlling pathogens that cause vascular wilt is not an easy task, the chemical fungicides that must be applied in the soil around the plant are ineffective especially for fungi such as *F. oxysporum*, which presents resistance structures that survive for long periods in the soil even in the absence of a host plant (Bani *et al.*, 2018; Chávez *et al.*, 2020; Srinivas *et al.*, 2019).

Regarding resistance, Pulido (2010), Manon *et al.* (2018), Carmona *et al.* (2020), and Leitão *et al.* (2020) state that this measure is the most effective and economically profitable for vascular fusarium in the field. Muñoz *et al.* (2019), Lamo & Takken (2020), and Leitão *et al.* (2020) add that the search for genotypes with different degrees of resistance allows a considerable decrease in the frequency of pesticide application, which minimizes the effects on human and environmental health, as well as a direct reduction in production costs. When trying to plan and apply new disease management methods, the objective should be a rational, effective, and safe control at a minimum cost as it is the use of resistant cultivars, many severe fungal diseases, as well as vascular wilt in economically important crops are treated in this way.

**Vascular discoloration.** Taking into account the scale (Table 3) proposed by Garcés *et al.* (2017), it was clearly evident that the vascular bundles of genotype 12U374 presented the most severe discoloration in the upper, middle, and lower part, and a reddish brown tone towards the interior of the three portions evaluated corresponding to different heights from the stem to the root (high, middle and low), allowing it to be

considered one of the most liable to attack by *FoPh* (Figure 3), even when compared to the commercial control, which showed discoloration grade 2 (intermediate). Most of the other materials showed intermediate discoloration grade except for 09U086, 09U089, 09U128, 12U357, and UN14, which showed intermediate discoloration grade. 12U357 and UN14 also revealed intermediate necrosis but only in the root portion, in the middle and upper part it was slight. Peru, on the other hand, showed slight discoloration in the three portions evaluated, and the genotypes 09U138 and 12U399 did not indicate vascular necrosis (Figure 3 and Figure 4A: SH=H, SM=M, SL=L); furthermore, the plants did not exhibit external symptoms such as generalized wilting, leaflet loss, growth reduction, or progressive drying, so they could be considered resistant to attack by this strain of *Fusarium oxysporum*.

In advanced stages, the roots of the plants show vascular discoloration, necrosis in the stalk base and stem, and discoloration of vascular bundles in addition to necrosis with a pattern of advancing towards the pith (Maurya *et al.*, 2019; Mendoza *et al.*, 2019). Cardona & Castaño (2019), in *Solanum lycopersicum* describe cross sections with necrotic processes, vascular tissue of dark brown color, being more noticeable at the junction point of the petiole with the stem, similar symptomatology to that presented in genotypes with an intermediate and severe degree of discoloration. In materials such as 12U374, externally, loss of the primary root and grayish-brown lesions at the point of emergence of lateral roots could be observed, as well as adventitious roots above the stem lesion (Figure 4B).



**Figure 3.** Degree of vascular discoloration of the 40 Goldenberry genotypes in the (H), middle (M), and low (L) of the plant.

Berruezo (2018) and Patiño (2020) state that vascular necrosis is due to the fact that xylem tracheids become obstructed, preventing communication between the cells of the conducting vessels and the transport of substances, which generates pronounced necrosis; what was observed in genotypes

09U086, 09U089, 09U128, 12U357, and UN14, suggests that it was difficult for the pathogen to colonize the stem of these genotypes, and for this reason its advance occurred only in the area near the inoculation point, thus confirming the ability of this microorganism to mobilize via the vascular system.




**Figure 4.** Cross section of Goldenberry stem. **A.** Healthy plant without vascular discoloration **B.** Plant with discoloration, caused by *Fusarium oxysporum* f. sp. *Physali*.

**Susceptibility index (SSI).** To determine the effect of *Fusarium oxysporum* f. sp. *physali* (*FoPh*) on the forty genotypes the susceptibility index was calculated, which made it possible to identify the outstanding materials considering the traits evaluated (plant height, AUDPC, incidence, and vascular discoloration), resulting in a relative value, by which the genotypes could be compared in general. In the analysis carried out, the response of the genotypes was separated into four groups: Very resistant (VR), Resistant (R), Susceptible (S), and Very susceptible (VS).

Figure 5 shows that many of the genotypes evaluated were in the very susceptible

category (VS), far from the materials 12U399 and 09U138 which were classified as very resistant (VR), since their susceptibility index was 0; on the other hand, the commercial control reached an SSI of 1 being one of the most susceptible (VS), the above, coincide with that reported by Osorio *et al* (2017) who by conglomerate analysis classified 09U138 as (VR), this source resistance to *FoPh* is apparently related with wild-type germplasm. Mayorga *et al.* (2019) in field tests and soils with high inoculum pressure observed an opposite reaction in which the material 09U138 was (S) to *FoPh*.



**Figure 5.** Classification of Goldenberry genotypes according to the susceptibility index MR (Very Resistant), R (Resistant), S (Susceptible), and MS (Very Susceptible). The values were normalized in relation to the commercial control.

The results obtained for 12U399 and 09U138 showed good potential to be incorporated in breeding programs to develop genotypes resistant to *FoPh* populations. Peru showed a certain degree of tolerance to the fungus indicating an index of 0.5, genotype that together with 12U399 and 09U138 should be evaluated in other regions and against other strains of the pathogen. UN34, 13U407, 09U089, Sylvania, UN26, and UN52 indicated an ISS that ranged between 0.83 and 0.96, and that are therefore considered susceptible (S), i.e., *F. oxysporum* in these plants is capable of penetrating, infecting, and causing symptoms characteristic of the disease and even causing death, which in the field could generate total losses to the farmer. The rest of the genetic materials, as well as the commercial control, are considered very susceptible (VS), although the highest SSI was for UN35 with 1.57. Finally, it should be remembered that

the use of resistant genotypes is one of the most appropriate strategies to reduce losses caused by the disease, since production costs are not increased, and the annual application of chemical products is reduced. The resistant genotypes drastically reduce environmental contamination, health, and food risks, and it also guarantees the success of other management tactics.

## CONCLUSIONS

Genotypes 09U138 and 12U399 from the first to the last evaluation did not show symptoms such as basal leaf wilt, loss of turgor, epinasty, chlorosis, stem prostration, or necrosis of the stem or roots, so they can be considered resistant to the attack of this strain of *F. oxysporum* f. sp. *physali*, however, the results obtained are an advance in the

search for resistance of this pathogen and the genotypes should be evaluated in the field, with the same isolation and with other pathogenic strains of the fungus.

Genotypes 09U138 and 12U399 could be used in breeding programs given their resistance to such an important disease as vascular wilt. Although the resistance to *Fusarium* shown by the other materials did not meet the expectations of this work, it is likely that these genotypes have characteristics of importance for the agro-industrial sector, where their potential should be considered.

### ACKNOWLEDGMENTS

The authors express their gratitude to the Universidad de Nariño, Gobernación de Nariño, and the Sistema General de Regalías, agreement 1509-14, for financing this study. Also, to Agrosavia C.I. Obonuco, especially to David Rodríguez and María Camila Ortega for their constant support in the field and laboratory, and finally to Dr. Pedro Uribe Mejía for his valuable contributions to this document.

**Conflict of interest:** The authors declare that there is no conflict of interest.

### BIBLIOGRAPHIC REFERENCES

- Agencia UNAL. (2018). Hongos y bacterias controlarían plaga de uchuva. Retrieved from <http://agenciadenoticias.unal.edu.co/detalle/hongos-y-bacterias-controlarian-plaga-de-la-uchuva>
- Agrios, G. (2005). *Plant Pathology*. (5ed). Nueva York: Elsevier Academic Press. 922p.
- Agudelo, N. (2020). Evaluación de la actividad antifúngica de ditiocarbamatos de alquilo derivados de vinilcetonas contra *Fusarium oxysporum* en plantas de uchuva (*Physalis peruviana*) bajo condiciones de invernadero. Retrieved from <https://repository.unimilitar.edu.co/handle/10654/3813>
- Alvarado, P. (2005). Identificación de cepas patógenas de *Fusarium* Link. causantes de nueva patología en el cultivo de calas de colores bajo condiciones productivas de invernadero en Chile. Retrieved from <https://biblioteca.inia.cl/handle/123456789/36791>
- Alves, D.; Simões, R.; Laurindo, B.; Laurindo, R.; Fonseca, F.; Damião, C.; Nick, C.; da Silva, H. (2017). Artificial neural network for prediction of the area under the disease progress curve of tomato late blight. *Scientia Agrícola*. 4(1): 51-59. doi: <https://doi.org/10.1590/1678-992X-2015-0309>
- Ángel, C.; Robledo, J.; Castaño, J. (2018). Comparación de métodos de inoculación de *Fusarium solani* f. sp. *passiflorae* en plántulas de maracuyá (*Passiflora edulis* f. *flavicarpa*). *Rev. U.D.C.A Act. & Div. Cient.* 21(1): 23-31. doi: <https://doi.org/10.31910/rudca.v21.n1.2018.659>
- Arellano, M. (2018). Detección de *Fusarium oxysporum* en cultivos de uvilla (*Physalis peruviana* L.) en la Sierra norte y centro del Ecuador. Retrieved from <http://repositorio.puce.edu.ec/handle/22000/14683>
- Bani, M.; Pérez, A.; Rubiales, D.; Rispaill, N. (2018). Physical and Chemical Barriers in Root Tissues Contribute to Quantitative Resistance to *Fusarium oxysporum* f. sp. *pisi* in Pea. *Frontiers in plant science*. 9(199): 1-16. doi: <https://doi.org/10.3389/fpls.2018.00199>
- Berrueto, L. (2018). Caracterización morfológica, biológica y molecular de los complejos *Fusarium oxysporum* y *Fusarium solani* asociados al cultivo de *Nicotiana tabacum* L. en el Noroeste Argentino. Retrieved from <https://ri.conicet.gov.ar/handle/11336/80238>
- Bocianowski, J.; Tratwal, A.; Kamila N. (2020). Genotype by environment interaction for área under the disease-progress curve (AUDPC) value in spring barley using additive main effects and multiplicative interaction model. *Australasian Plant Pathology*. 49: 525-529. doi: <https://doi.org/10.1007/s13313-020-00723-7>



- Burbano, O. (2020). Resistencia de plantas a patógenos: una revisión sobre los conceptos de resistencia vertical y horizontal. *Revista Argentina de Microbiología*. 52(3): 245-255. doi: <https://doi.org/10.1016/j.ram.2020.04.006>
- Cardona, L.; Castaño, J. (2019). Comparación de métodos de inoculación de *Fusarium oxysporum* f. sp. *lycopersici* (Sacc.) Snyder & Hansen, causante del marchitamiento vascular del tomate. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales*. 43(167): 227-233. doi: <https://doi.org/10.18257/raccefyn.854>
- Carmona, S.; Burbano, D.; Gómez, M.; López, W.; Ceballos, N.; Castaño, J.; Simbaqueda, J.; Soto, M. (2020). Characterization of pathogenic and nonpathogenic *Fusarium oxysporum* Isolates Associated with commercial tomato crops in the Andean Region of Colombia. *Pathogens*. 9(70): 1-23. doi: <https://doi.org/10.3390/pathogens901007>
- Castellanos, L.; Torrado, J.; Céspedes, N. (2021). Alternativas biológicas para el control de *Fusarium oxysporum* en arveja en Pamplona, Norte de Santander. *Revista de investigación Agraria y Ambiental*. 12: 1-13. doi: <https://doi.org/10.22490/21456453.3650>
- Castro, F.; Gentzel, I.; Mackey, D.; Alonso, A. (2020). Metabolomics as an emerging tool for the study of Plant-Pathogen Interactions. *Metabolites*. 10(52): 1-23. doi: <https://doi.org/10.3390/metabo10020052>
- Cubedo, L. (2008). Manejo biorracional de la pudrición de la corona y raíz (*Fusarium oxysporum* f. sp. *Radicallycopersici*) del tomate (*Lycopersicon esculentum* Mill.) Retrieved from <https://tesis.ipn.mx/handle/123456789/3806>
- Chañag, H.; Viveros, J.; Álvarez, S.; Criollo, H.; Lagos, L. (2017). Evaluation of tree tomato genotypes *Cyphomandra betacea* (Cav.) Sendt. for resistance to *Phytophthora infestans* Mont. de Bary sensu lato attacks. *Revista Colombiana De Ciencias Hortícolas*. 11(1): 11-20. doi: <https://doi.org/10.17584/rcch.2017v11i1.4725>
- Chávez, C.; Gómez, S.; Restrepo, H. (2019). Physiological, biochemical and chlorophyll fluorescence parameters of *Physalis peruviana* L. seedlings exposed to different short-term waterlogging periods and *Fusarium* Wilt Infection. *Agronomy*. (5):20. doi: <https://doi.org/10.3390/agronomy9050213>
- Chávez, C.; Gómez, S.; Restrepo, H. (2020). Physiological responses to the foliar application of synthetic resistance elicitors in cape gooseberry seedlings infected with *Fusarium oxysporum* f. sp. *physali*. *Plants* 9(2): 24. doi: <https://doi.org/10.3390/plants902012a>
- Díaz, J.; Acuña, W.; Yáñez, J. (2019). Diversidad de comunidades fúngicas que habitan en tejido foliar sintomático de la uvilla *Physalis peruviana* en los Andes Ecuatorianos. *Revista ecuatoriana de Medicina y Ciencias Biológicas*. 40(2): 77-86. doi: <https://doi.org/10.26807/remcb.v40i2>
- EVA-Evaluaciones Agropecuarias Municipales. (2021). Cultivo de uchuva. Retrieved from <https://experience.arcgis.com/experience/17859d5712b046fca6b0df5781e0b560/page/EVA-Departamental/>
- Gayosso, O.; López, A.; Marroquín, J.; López, K.; Hidalgo, D.; Chávez, G. (2021). Evaluación de la respuesta de diferentes genotipos de tomate a *Fusarium oxysporum* raza 3. *Revista Mexicana de Ciencias Agrícolas*. 12(3): 409-420. doi: <https://doi.org/10.29312/remexca.v12i3.2284>
- Garcés, F.; de Borda, C.; Schmidt, E.; Bouzon Z.; Stadnik, M. (2017). Delayed upward colonization of xylem vessels is associated with resistance of common bean to *Fusarium oxysporum* f. sp. phaseol. *European Journal of Plant Pathology*. 149: 477-489. doi: <https://doi.org/10.1007/s10658-017-1197-6>
- García, D.; Dávila, L.; Gonzáles, A.; Caro, A.; Cortes, A. (2021). Sistema gnotobiótico para seleccionar microorganismos con potencial de biocontrol contra *Fusarium oxysporum* f. sp. *Physali*. *Revista Colombiana de Ciencias Hortícolas*. 15(1): e11699. doi: <https://doi.org/10.17584/rcch.2021v15i1.11699>
- Giraldo, C.; Velandia, D.; Fisher, G.; Gómez, S.; Martínez, L. (2020). Respuesta hiperespectral de plantas de uchuva (*Physalis peruviana* L.), inoculadas con *Fusarium oxysporum* f. sp. *physali*, para la detección del marchitamiento vascular. *Revista Colombiana de Ciencias Hortícolas*. 14(3): 301-313. doi: <https://doi.org/10.17584/rcch.2020v14i3.10938>

- González, A. (2019). Hongos asociados a la pudrición del cacao *Theobroma cacao* L. en el soconusco, Chiapas y patogenicidad y control *in vitro* de *Nodulosporium*. Retrieved from <http://repositorio.uaaan.mx:8080/xmlui/handle/123456789/46232>
- Gordon, T. (2017). *Fusarium oxysporum* and the *Fusarium* wilt syndrome. *Annual Review Phytopathol.* 55(1): 23–39. doi: <https://doi.org/10.1146/annurev-phyto-080615-095919>
- Insuasty, C.; Acosta, J.; Salazar, C.; Betancourth, C. (2014). Evaluación de cepas de *Trichoderma* spp. para el manejo del amarillamiento de arveja causado por *Fusarium oxysporum*. *Corpoica. Ciencia y Tecnología Agropecuaria.* 15(2): 237-249.
- Islas, C. (2021). Estudio de efecto antagónico de una cepa bacteriana contra *Fusarium oxysporum* en jitomate *Solanum lycopersicum* L. Retrieved from <https://repositorioinstitucional.buap.mx/handle/20.500.12371/11786>
- Joshi, R. (2018). A review of *Fusarium oxysporum* on its plant interaction and industrial use. *Journal of Medicinal Plants Studies.* 6(3): 112-115. doi: <http://dx.doi.org/10.22271/plants.2018.v6.i3b.07>
- Kaur, B.; Bathia, D.; Mavi, S. (2021). Eighty years of gene-for-gene relationship and its applications in identification and utilization of R genes. *Journal of Genetics.* 100(50): 1-17. doi: <https://doi.org/10.1007/s12041-021-01300-7>
- Lamo, F.; Takken F. (2020). Biocontrol by *Fusarium oxysporum* Using Endophyte-Mediated Resistance. *Frontiers in plant Science.* 11(37): 1-15. doi: <https://doi.org/10.3389/fpls.2020.00037>
- Leitão, S.; Malosetti, M.; Song, O.; Eeuwijk, F.; Rubiales, D.; Vaz Patto, C. (2020). Natural Variation in Portuguese Common Bean Germplasm Reveals New Sources of Resistance Against *Fusarium oxysporum* f. sp. phaseoli and Resistance-Associated Candidate Genes. *Phytopathology.* 110:633-647. doi: <https://doi.org/10.1094/PHTO-06-19-0207-R>
- Manon, M.; Richard, S.; Gratias, A.; Meyers, B.; Geffroy, V. (2018). Molecular mechanisms that limit the costs of NLR-mediated resistance in plants. *Molecular Plant Pathology.* 9(11): 2516-2523. doi: <https://doi.org/10.1111/mpp.12723>
- Marín, J.; Hoyos, L.; Botero, M. (2018). Detección de plantas asintomáticas de *Solanum lycopersicum* L. infectadas con *Fusarium oxysporum* usando espectroscopia de reflectancia VIS. *Revista Colombiana de Ciencias Hortícolas.* 12(2): 436-446. doi: <https://doi.org/10.17584/rcch.2018v12i2.7293>
- Maurya, S.; Dubey, S.; Kumari, R.; Verma, R. (2019). Management tactics for fusarium wilt of tomato caused by *Fusarium oxysporum* f. sp. *lycopersici* (Sacc.): A review. *International Journal of Research in Pharmacy and Pharmaceutical Sciences.* 4(5): 1-7.
- Mayorga, F.; Arguelles, J.; Rodríguez, E.; González, C.; Ariza, C.; Barrero, L. (2019). Producción y calidad fisicoquímica del fruto de *Physalis peruviana* L. en relación con respuesta de resistencia a *Fusarium oxysporum* f. sp. *Physali.* *Agronomía Colombiana.* 37(2): 120-128. doi: <https://doi.org/10.15446/agron.colomb.v37n2.77550>
- Mendoza, D.; López, C.; Valdés, T.; Velásquez, T.; López G.; Martínez, R. (2019). Relación entre la nutrición en tomate (*Solanum lycopersicum*) y la severidad de la fusariosis. Retrieved from <https://www.omniascience.com/books/index.php/proceedings/catalog/view/106/448/886-1>
- Moreno, C.; Izquierdo, L.; Ongena, M.; Kloepper, Cotes, A. (2019). Soil sterilization, pathogen and antagonist concentration affect biological control of *Fusarium* wilt of cape gooseberry by *Bacillus velezensis* Bs006. *Plant soil.* 435: 39-55. doi: <https://doi.org/10.1007/s11104-018-3866-4>
- Muñoz, M.; Acuña, I.; Sagredo, B. (2019). Resistencia varietal al tizón tardío de la papa. En: Acuña, I.; & Bravo, R. (eds.). *Tizón tardío de la papa: Estrategias de manejo integrado con alertas temprana.* pp. 100-117. Chile: Instituto de investigaciones Agropecuarias - INIA. 399p.
- Osorio, J.; Enciso, F.; González, C.; Pozo, N.; Mueller, L.; Barrero, L. (2016). Association analysis for disease resistance to *Fusarium oxysporum* in cape gooseberry (*Physalis peruviana* L). *BMC Genomics.* 17(248): 1-16. doi: [doi:10.1186/s12864-016-2568-7](https://doi.org/10.1186/s12864-016-2568-7)
- Patiño, M. (2020). Evaluación de la resistencia de introducciones de *Passifloras* spp. a la “secadera”

- causada por *Fusarium* spp. de zonas productoras. Retriev from <https://repositorio.unal.edu.co/handle/unal/78567>
- Pouralibaba, R.; Pérez-De-Luque, A.; Rubiales, D. (2017). Histopathology of the infection on resistant and susceptible lentil accessions by two contrasting pathotypes of *Fusarium oxysporum* f.sp. lentis. *European Journal Plant Pathol.* 148: 53-63. doi: <https://doi.org/10.1007/s10658-016-1068-6>
- Pulido, V. (2010). Evaluación de la resistencia y susceptibilidad de accesiones élite de germoplasma de uchuva (*Physalis peruviana* L.) al hongo *Fusarium oxysporum* schldt. Retriev from <https://www.grin.com/document/280811>
- Pulido, V.; Rodríguez, E.; Barrero, L.; Gonzáles, C. (2011). Búsqueda de la resistencia a la marchitez vascular en germoplasma de uchuva y especies relacionadas. En: Gonzáles, C.; Barrero, L. (eds.). *Estudio de la marchitez vascular de la uchuva para el mejoramiento genético del cultivo*. pp: 21-29. Bogotá: Kimpres.
- Rodríguez, E. (2013). Caracterización de aislamientos de *Fusarium* spp. obtenidos de zonas productoras de uchuva (*Physalis peruviana*) en Cundinamarca y Boyacá. Bogotá, Colombia: Facultad de Agronomía, Universidad Nacional de Colombia. 99p.
- Rodríguez, A.; Pedraza, L. (2019). Efecto del método de inoculación y la presencia de heridas en la raíz sobre el marchitamiento vascular de plantas de uchuva (*Physalis peruviana*) causado por *Fusarium oxysporum*. Retriev from <https://n9.cl/ewnmz>
- Rodríguez, A. (2020). Compatibilidad de fungicidas químicos, biológicos y de origen vegetal sobre el hongo benéfico *Trichoderma harzianum*, controlador de *Fusarium oxysporum* en plantas de tomate (*Solanum lycopersium*). Retriev from <http://hdl.handle.net/20.500.12010/8628>
- Rose, S.; Parker, M.; Punja, Z. (2003). Efficacy of biological and chemical treatments for control of *Fusarium* root and stem rot on greenhouse cucumber. *Plant Disease.* 87: 1462-1470. doi: <https://doi.org/10.1094/PDIS.2003.87.12.1462>
- Ruiz, L.; Castellanos, L.; Villamizar, C. (2018). El cultivo de la uchuva (*Physalis peruviana*). *Revista Científica Agroecosistemas.* 6(1): 46-53.
- Sánchez, D.; Hernández, R.; Andrade, M.; López, V.; Tejacal, A.; Juárez, P. (2017). Eficacia de seis fungicidas en el control de *Mycosphaerella citri* Whiteside en naranja, Var. 'Valencia' en Tlayecac, Morelos, México. *Revista Centro Agrícola.* 44(3): 71-79.
- Simbaqueba, J.; Catanzariti, C.; González, D.; Jones. (2018). Evidence for horizontal gene transfer and separation of effector recognition from effector function revealed by analysis of effector genes shared between cape gooseberry and tomato- infecting formae speciales of *Fusarium oxysporum*. *Molecular Plant Pathology.* 19: 2302-2318. doi: [10.1111/mp.12700](https://doi.org/10.1111/mp.12700)
- Simbaqueba, J. A.; Rodríguez, E.; Burbano, D.; Gonzáles, C.; Caro, A. (2021). Putative Novel Effector Genes Revealed by the Genomic Analysis of the Phytopathogenic Fungus *Fusarium oxysporum* f. sp. *physali* (Foph) That Infects Cape Gooseberry Plants. 11: 1-15. doi: <https://doi.org/10.3389/fmicb.2020.593915>
- Srinivas, C.; Nirmala, D.; Narasimha, K.; Dhananjaya, C.; Lakshmeesha, T.; Singh, B.; Kumar, M.; Niranjana, S.; Hashem, A.; Abdulaziz, A.; Tabassum, B.; Abd\_Allahg, E.; Nayaka, C.; Srivastava, R. (2019). *Fusarium oxysporum* f. sp. *lycopersici* causal agent of vascular wilt disease of tomato: Biology to diversity- A review. *Saudi Journal of Biological Sciences.* 26(7): 1315-1324. doi: <https://doi.org/10.1016/j.sjbs.2019.06.002>
- Valderrama, D. (2018). Control del marchitamiento vascular de la uchuva basado en mezclas de microorganismos rizosféricos provenientes de suelos potencialmente supresivos. Retriev from <https://repositorio.unal.edu.co/handle/unal/64732>
- Vásquez, L.; Castaño, J. (2017). Manejo integrado de la marchitez vascular del tomate [*Fusarium oxysporum* f. sp. *Lycopersici* (SACC.) W.C. Snyder & H.N. Hansen]: una revisión. *Revista U.D.C.A.* 20(2): 363-374. doi: <https://doi.org/10.31910/rudca.01234226>
- Wang, B.; Jeffers, S. (2002). Effects of cultural practices and temperature on *Fusarium* root and crown rot of container-grown hostas. *Plant Disease.* 86: 225-231. doi: <https://doi.org/10.1094/PDIS.2002.86.3.225>