



Determinants of nitrogen use efficiency in coffee crops. A review

Determinantes de la eficiencia en el uso del nitrógeno en los cultivos de café. Una revisión

Siavosh Sadeghian-Khalajabadi¹; José-Ricardo Acuña-Zornosa²; Luis-Fernando Salazar-Gutiérrez³; Juan-Camilo Rey-Sandoval⁴

ARTICLE DATA

1. Scientific Researcher III, Ph.D. Centro Nacional de Investigación del Café - Cenicafé, Manizales, Colombia, siavosh.sadeghian@cafedecolombia.com
<https://orcid.org/0000-0003-1266-0885>
2. Scientific Researcher III, Ph.D. Centro Nacional de Investigación del Café - Cenicafé, Manizales, Colombia, ricardo.acuna@cafedecolombia.com
<https://orcid.org/0000-0001-6935-2264>
3. Scientific Researcher II, Ph.D. Centro Nacional de Investigación del Café - Cenicafé, Manizales, Colombia, luisfernando.salazar@cafedecolombia.com
<https://orcid.org/0000-0003-2302-4825>
4. Research Assistant, M.Sc. Centro Nacional de Investigación del Café - Cenicafé, Manizales, Colombia, juan.rey@cafedecolombia.com
<https://orcid.org/0000-0003-0040-3291>

Cite: Sadeghian-Khalajabadi, S.; Acuña-Zornosa, J.-R.; Salazar-Gutiérrez, L.-F.; Rey-Sandoval, J.-C. (2022). Determinants of nitrogen use efficiency in coffee crops. A review. *Revista de Ciencias Agrícolas*. 39(2): 198-219.
doi: <https://doi.org/10.22267/rcia.223902.183>

Received: August 28 2021.
Accepted: March 9 2022.



ABSTRACT

World production of coffee (*Coffea* spp.) has increased in recent decades due to its growing demand. Nitrogen (N) is one of the most required nutrients by coffee crops and in many cases, the most yield-limiting; nevertheless, a high percentage of supplied N is lost, generating pollution, greenhouse gases, and economic losses. Given this scenario, it is a priority to adopt practices that increase nitrogen use efficiency (NUE) related to the capacity of plants to absorb and use N to produce biomass. This article provides context about NUE and determining factors in coffee: plant, soil, climate, and management. On the plant side, despite advances in genetics and the results of genotype evaluation, it may be a decade or more before improved coffee varieties with high NUE become available or used in world coffee farming. Both the physical, chemical, and biological properties of the soil and the elements of the climate, mainly temperature, precipitation, and solar radiation, have an influence on coffee growth and many of the processes that intervene in the N cycle and NUE; however, some components of these two factors are not fully controllable. Management of the factors that determine NUE should focus on reducing N losses in the system, increasing its uptake and utilization by plants, and maintaining or increasing productivity.

Keywords: *Coffea* spp.; climate; crop management; crop production; environment; plant nutrition; soil.

RESUMEN

La producción mundial de café (*Coffea* spp.) ha aumentado en las últimas décadas, como consecuencia de su creciente demanda. El nitrógeno (N) es uno de los nutrientes más requeridos por el cultivo del café y en muchas ocasiones el más limitante de la producción; sin embargo, un alto porcentaje de la cantidad suministrada de N se pierde, generando contaminación ambiental, gases de efecto invernadero y pérdidas económicas. Ante este panorama, se hace prioritaria la adopción de prácticas para aumentar la eficiencia de uso

del N (EUN), la cual se relaciona con la capacidad de las plantas para absorber y utilizar N con el propósito de producir biomasa. Este artículo provee contexto acerca de la EUN y los factores que la determinan en café: planta, suelo, clima y manejo. En cuanto a la planta, a pesar de los avances en genética y los resultados de la evaluación de genotipos, es posible que se pase una década o más, antes que las variedades mejoradas de café con alta EUN estén disponibles o se utilicen en la caficultura mundial. Tanto las propiedades físicas, químicas y biológicas del suelo como los elementos del clima, principalmente temperatura, precipitación y radiación solar, afectan el crecimiento de café y muchos de los procesos que intervienen en el ciclo del N y afectan la EUN; sin embargo, algunos componentes de estos dos factores no son totalmente controlables. El manejo de los factores que determinan la EUN debe enfocarse en reducir las pérdidas de N en el sistema, aumentar su absorción y utilización por las plantas y mantener o aumentar la productividad.

Palabras clave: *Coffea* spp.; clima; manejo del cultivo; producción de cultivos; medio ambiente; nutrición de las plantas; suelo.

INTRODUCTION

During the last three decades, coffee (*Coffea* spp.) production has increased steadily; it went from 93 million 60.0kg bags in 1990 to 165 million bags in 2019. Although coffee is grown today in about 60 countries, 95% of the total volume is produced in 16 of them and 62% in only three: Brazil (35%), Vietnam (18%), and Colombia (9%) (International Coffee Organization, 2021).

There are two ways to satisfy the growing demand for coffee beans; one through incorporating new areas of agricultural vocation, which are becoming increasingly smaller, and the other; through increased productivity on lands currently cultivated (Sadeghian, 2014). According to the above, the production goals proposed in the medium and long term depend on the efficient and effective management of nutrients to achieve the expected economic, social, and environmental benefits (Fixen *et al.*, 2015).

The continued extraction of nutrients by the yield, without the respective return, leads to the depletion of soil reserves and the subsequent reduction in production; conversely, the excessive use of fertilizers and unbalanced nutrition also generates

contamination with possible yield consequences (Moody & Bruulsema, 2020). In this sense, nitrogen (N) constitutes one of the two macronutrients with the highest demand and influence on crop production, among which coffee is no exception; hence, it is included in fertilization plans at relatively high rates (Raij *et al.*, 1996; Guimarães *et al.*, 1999; Sanzonowicz *et al.*, 2000; Sadeghian, 2013). Despite the above, more than half of the applied N can be lost in the environment (Lassaletta *et al.*, 2014), wasting the resource, increasing production costs and causing threats to the air, water, soil, and biodiversity, and emitting greenhouse gases, a situation that has increasingly questioned their use in recent years (Dobermann, 2005; Lassaletta *et al.*, 2014; World Meteorological Organization, 2019). The analysis by Lassaletta *et al.* (2014), based on the information collected by FAO during the last 50 years, revealed, for example, that in Greece, France, and the Netherlands, the nitrogen use efficiency (NUE) has increased in terms of yield of crops and total N inputs. While in India, Egypt, and China, increased fertilization has produced few agronomic benefits and significant environmental losses.

At the same time, large amounts of fertile soil rich in organic carbon (OC) and N are

constantly being lost, generating more greenhouse gases, mainly CO₂, CH₄, and N₂O (Lal, 2020), as a consequence of erosion and inappropriate management practices, a situation that affects root growth and increases the need to apply more N to sustain or improve yield. The described panorama turns on the alarms to conserve the soil and seek strategies that contribute to the efficient use of N. At the same time, alternatives are evaluated to increase yield and reduce environmental impacts (Batabyal, 2017). These strategies must integrate nitrogen fertilization with cultivars that are more efficient in their use and other cultural practices within good management practices (Sarkar & Baishya, 2017).

This article reviews the factors that determine NUE in coffee: plant, soil, climate, and management.

Definitions of nutrient use efficiency and NUE.

To refer to NUE, it is convenient to previously define the concept of nutrient use efficiency (NUTUE), which is one of the leading indicators of mineral nutrition concerning the processes by which plants acquire, transport, store, and use nutrients to produce biomass when low or high rates of nutrients are applied through fertilization (Mathur & Goel, 2017).

There are different definitions of NUTUE; however, most authors agree on referring to the yield or nutrient absorbed/removed per unit of nutrient or fertilizer applied (Bruulsema *et al.*, 2008), a statement comprising two components: nutrient absorption efficiency (the plant's ability to extract nutrients from the soil) and utilization efficiency (the ability to use nutrients to produce biomass). The

evaluation of the NUTUE is helpful in the differentiation of soils for their ability to supply nutrients to plants and differentiate species and cultivars by their ability to absorb and use nutrients to produce maximum yields (Sarkar & Baishya, 2017).

In agronomic research, the following indexes are commonly used to evaluate the efficiency of the applied nutrients: partial factor productivity, partial nutrient balance, agronomic efficiency, apparent nutrient recovery efficiency, internal utilization efficiency, and physiological efficiency (Table 1). It is essential to consider that the NUTUE changes in space and time; therefore, the indicators make more sense when obtained for particular conditions of the cultivation system, soil, climate, and management contexts (Sarkar & Baishya, 2017).

Based on the NUTUE concept, NUE can be defined as the yield obtained or N absorbed/removed per unit of N or nitrogen fertilizer applied. The two components of NUE would then be the efficiency or capacity of the plant to absorb N from the soil and the efficiency or ability to use N to produce biomass (Moll *et al.*, 1982; Hawkesford *et al.*, 2012). In this regard, it is necessary to consider that a part of the N that enters the system is removed from the field in the form of harvested biomass. At the same time, the amount contained in crop residues is incorporated into the soil and recycled, making part of the organic matter (OM) and inorganic N reserves. However, another component, called reactive N, is not recovered and is lost from the system in inorganic reduced forms (e.g., NH₃ and NH₄⁺), oxidized inorganic forms (e.g., NO_x, HNO₃, N₂O, and NO₃⁻), and organic compounds (e.g., urea, amines, and proteins) (Cassman *et al.*, 2002).

Table 1. Summary of commonly used concepts of nutrient use efficiency.

Definition	Formula
Partial factor productivity (PFP) is units of crop yield per unit of applied nutrient.	$PFP = Y_F/F$ or $PFP = (Y_0/F) + AE$
Partial nutrient balance (PNB) is units of nutrient uptake per unit of applied nutrient.	$PNB = U_F/F$
Agronomic efficiency (AE) of the applied nutrient is units of crop yield increase per unit of applied nutrient.	$AE = (Y_F - Y_0)/F$ or $AE = ANR \times PE$
Apparent nutrient recovery efficiency (ANR) is an increase in units of nutrient uptake per unit of applied nutrient.	$ANR = (U_F - U_0)/F$
Internal utilization efficiency (IE) of a nutrient is units of crop yield per unit of nutrient uptake.	$IE = Y_F/U_F$
Physiological efficiency (PE) of the applied nutrient is an increase in units of crop yield per unit increase in nutrient uptake from fertilizer.	$PE = (Y_F - Y_0)/(U_F - U_0)$

F, amount of nutrient applied (kg ha^{-1}); Y_F and Y_0 , crop yield with and without applied nutrient (kg ha^{-1}), respectively; U_F and U_0 , total nutrient uptake in aboveground biomass at maturity (kg ha^{-1}) in fertilized and unfertilized control plot, respectively. Original information from Cassman *et al.* (2002) and Dobermann (2007), summarized by Sarkar & Baishya (2017).

The EU Nitrogen Expert Panel (EUNEP, 2015) sustains that there is currently no uniform and robust methodology and protocol for estimating NUE. A more accessible indicator is then proposed, applicable to agriculture and production systems and food consumption based on the mass balance principle, using input and output data of N for its calculation: $NUE = N \text{ output}/N \text{ input}$. The NUE values must be interpreted as productivity (N output) and N surplus (the difference between N input and N output harvested).

It is possible to divide NUE studies into two groups: i) researches that generate indexes for one or more crops at specific sites for short periods (usually less than two years), using mainly primary information; for example, Fageria (2014) in cereals and bean and ii) comparative studies that use secondary information to evaluate regional changes (countries and continents) of some indices over more extended periods (more

than ten years); for example, the reports for cereals and soybean of Cassman *et al.* (2002), Dobermann & Cassman (2002), Lassaletta *et al.* (2014), EUNEP (2015) and Fixen *et al.* (2015).

Determinants of NUE in coffee crops. N is one of the most required nutrients by the coffee crop (Martinez *et al.*, 2020; Cenicafé, 2020), and, in many occasions, N is the most limiting yield (Sadeghian, 2008; Sadeghian, 2009). Fertilizers that contain N that contribute significantly to crop yield (Fageria, 2014). However, a high percentage of the quantity of N supplied is lost (more than 40%) (Prezotti *et al.*, 2000; Bote *et al.*, 2018b; Bess, 2020), which generates environmental pollution, greenhouse gases, and economic losses; therefore, adopting practices to increase NUE becomes a priority. This objective is not easy to achieve because the N cycle is complex. N is easily lost from agriculture to the environment (EUNEP, 2015).

The same factors that determine crop productivity also influence NUTUE and, therefore, NUE: plant, soil, climate, and management.

Plant. Having efficient cultivars in the use of nutrients contributes to improving the profitability of agricultural activity and reduces the impact on the environment. There are different definitions for an efficient plant in using nutrients, as discussed by Fageria (2014). Soil Science Society of America (2008) defines a nutrient-efficient plant as: "A plant that absorbs, translocates, or uses more of a specific nutrient than another plant under relatively low nutrient availability conditions in the soil or growing media." Regarding N, Graham (1984) refers to the ability of a genotype to produce higher yields under conditions of low nutrient content in the soil compared to other genotypes. The traits that distinguish the efficient species/cultivars in N absorption are related to the morphology of the root and the activity of the transporter; in turn, NUE is a consequence of the metabolic processes that favor the plant's capacity to assimilate and remobilize N towards the organs of interest (Masclaux-Daubresse *et al.*, 2010); in the case of coffee, the bean contained in the fruits.

According to Engels *et al.* (2012), biomass partition between plant organs and their control mechanisms are crucial in crop production. The partitioning of N refers to the differential distribution of N among different sinks (Bhatla & Lal, 2018; Taiz *et al.*, 2021).

Coffee species and varieties differ in the uptake/absorption and partitioning of N; therefore, genetic variability in NUE is also expected, which can be helpful in crop breeding programs (Cardoso, 2010). For Arabica coffee (*Coffea arabica* L.), Malavolta *et al.* (2002), Cardoso (2010), Amaral *et al.* (2011a), Amaral *et al.* (2011b), Martins *et*

al. (2015), and Martinez *et al.* (2020) report differences in NUE between cultivars for Brazilian conditions. Similar results have been obtained for the Conilon coffee clones (*Coffea canephora* Pierre ex Froehner) by Prezotti & Bragança (2013), Colodetti *et al.* (2014), and Machado *et al.* (2016).

Martins *et al.* (2015) affirm the existence of genetic variability in coffee genotypes of the same species in terms of the absorption and nutrient use efficiency; nevertheless, they consider it necessary to better understand the absorption and utilization mechanisms of nutrients, as well as nature and genetic inheritance concerning efficiency. For these authors, Conilon coffee has a high NUE, which is suggested for selecting effective and responsive genotype indexes. For colombian conditions such differences have not been corroborated, as revealed by the research results carried out with *C. arabica* L. variety Castillo® under field conditions for 270 days after sowing (López, 2009), probably as a consequence of the small genetic amplitude among the evaluated accessions.

One of the difficulties in the selection of N-efficient genotypes in coffee is related to the relatively long cycle of the crop since the studies on perennial plants require more time to reach conclusive results, in contrast to the ones conducted on corn, wheat and rice. Researchers try to overcome this obstacle by evaluating materials in the seedling state, either using soil (Colodetti *et al.*, 2014; Machado *et al.*, 2016) or nutritive solutions (Cardoso, 2010; Martinez *et al.*, 2020) in the field for two years Amaral *et al.*, (2011a), Amaral *et al.*, (2011b). It is not the most desired procedure, because neither does it represent crop conditions nor the variations in yield exhibited by the plantations between one year to the next. N use efficient genotypes are not yet an option disseminated by researchers and adopted by coffee growers for the reasons above.

Improving NUE is a biological, agronomic, and environmental problem. Advances in understanding the functional biology of N response and NUE have revealed how N source and N concentration have a genome-wide and plant-wide impact, involving genes from multiple biological processes (Sinha *et al.*, 2020).

Several NUE definitions exist, but N uptake and utilization efficiency are the most relevant biological challenge (Mandal *et al.*, 2018). Identifying proper metabolic pathways to genetically improve the NUE of crops is hampered by incomplete characterization of plant phenotype and genotype. (Mandal *et al.*, 2018). Genetic improvement of NUE is often attempted within high-yielding varieties rather than searching for it within the genetic diversity of available germplasm (Mandal *et al.*, 2018; Sinha *et al.*, 2020).

Most of the current varieties accumulate important agronomical traits selected on fertile soils. The allelic diversity for the NUE traits would be shallow and even lack such alleles on nitrogen-limited soils (Sinha *et al.*, 2020). Above is the case for the evaluation of *C. arabica* genotypes for NUE. Research carried out with this species has been characterized by the use of genotypes obtained in genetic improvement programs aimed at developing coffee varieties with high productive yields in environments that are not deficient in N (López, 2009; Martins *et al.*, 2015). Therefore, it is crucial to identify contrasting genotypes or varieties so that germplasm can be ranked in increasing or decreasing order of NUE and thus make genetic associations using association genotyping techniques (Li *et al.*, 2017). Molecular markers for selection/improvement can also be developed using conventional germplasm screening or mining available genomic sequences (Sinha *et al.*, 2020).

N is mainly taken up through the roots and transported throughout the plant via the xylem. N compounds are also recycled and remobilized from internal stores or senescent tissues through the phloem to demand sites for bean filling, as is the coffee case (Filho & Malavolta, 2003). Genes involved in N uptake, assimilation, and remobilization processes are necessary to determine NUE, making NUE a complex quantitative trait (Mandal *et al.*, 2018).

On a whole-plant scale, the response to N encompasses other genes and biological processes that contribute to NUE, including its regulation of carbon metabolism, redox metabolism, and root architecture. This is how genes involved in N sensing and upstream processes, including epigenetic regulation involving miRNAs, have been identified (Li *et al.*, 2017).

Another major biological challenge for the genetic improvement of NUE is that neither the phenotypic traits nor the genetic alleles that determine it are clearly defined (Li *et al.*, 2017; Mandal *et al.*, 2018). Therefore, there is a great need to use chemical compound analogs from different N sources to carry out phenotypic screening for NUE-related genes/alleles (Mandal *et al.*, 2018).

Soil. The ability of the soil to supply N to plants and NUE depends on its physical, chemical, and biological properties, mainly OM, flora and fauna, pH, texture, topography, aeration, moisture, temperature, and cation exchange capacity/effective cation exchange capacity (CEC/ECEC). These intervene in N cycle processes (fixation, mineralization, immobilization, nitrification, denitrification, leaching, volatilization, absorption, and removal by yield) and, therefore, affect its use efficiency (Havlin *et al.*, 2017). There are countless reports in this regard, of which this item addresses some fundamental aspects.

In Figure 1, a conceptual model is presented, proposed by Balasubramanian *et al.*, (2004). This describes the three main components (demand, supply, and losses) that control the recovery efficiency of N applied in fertilization variables and key processes.

OM, particularly the stable fraction (humus), is the primary N source for plants (Pilbeam, 2015; Raji, 2011), including coffee (Sadeghian, 2011); therefore, the response of crops to N supply and NUE depends on the mineralization of OM and crop residues. Through the constant growth of the roots and the entry of plant remains and other organisms into the system, residues are permanently incorporated into the soil which replenishes the OM contents, avoiding the depletion of this resource due to mineralization over time (Weil & Brady,

2017). According to Cardona & Sadeghian (2005) reports, more than 4.0Mg ha⁻¹ of organic material annually enter the unshaded coffee monoculture and 8.0Mg ha⁻¹ in shaded plantations of *Inga* spp., amounts that come to contribute to the system 100 and 200kg ha⁻¹ of N, respectively. Recent reports suggest that the biomass of the vegetative organs of a variety Castillo® coffee crops in Colombia represent 58.0Mg ha⁻¹ at 60 months after sowing, reaching 855kg ha⁻¹ of N (Cenicafé, 2020). On the other hand, the loss of the superficial soil horizon due to water erosion reduces the OM, altering other properties such as bulk density, structural stability, and moisture retention, among others, which can minimize coffee productivity by up to 51.0% (Hincapié & Salazar, 2011).




Figure 1. Conceptual model representing the three main control boxes (e. g., nitrogen demand, supply, and losses) and their effective processes and variables regulating fertilizer N use efficiency (NUE). (Balasubramanian *et al.*, 2004).

Soil OM levels can be very contrasting in coffee-producing countries and regions, related to the origin of the soils and environmental conditions. In this regard, it is enough to mention the contents of less than 3.0% in Oxisols from Brazil compared to Andisols from Colombia with 20.0% or more (Sadeghian, 2011).

Due to the activity of heterotrophic organisms in the soil, a fraction of N contained in humus becomes available to plants by mineralizing OM. The magnitude and variations of these N contributions depend on the quantity and quality of OM and organic residues and the environmental factors that affect this process: temperature, moisture, aeration, and pH (Havlin *et al.*, 2017). As an indicator of mineralization, microbial activity may vary between sites over time and, eventually, be higher in unshaded plantations than in shaded ones due to their higher temperature (Cardona and Sadeghian, 2005). This activity can intensify when the pH increases by liming, increasing nitrate levels resulting from more significant mineralization of the soil OM (Sadeghian & Díaz-Marín, 2020). According to Cannavo *et al.* (2013), the amount of mineralized N in the first 10.0cm of the profile of an Andisol represents up to 200kg ha⁻¹ year⁻¹ of N in coffee plantations under shade, reaching a large part of the crop's requirements. In turn, Babbar & Zak (1994) corroborated that OM mineralization depends on the coffee production system and changes over time, influenced by periods of rain and drought. The results obtained by these authors for conditions in Costa Rica reveal amounts equivalent to 148kg ha⁻¹ year⁻¹ in coffee crops under shade and 110kg ha⁻¹ year⁻¹ in unshaded plantations. Measurements made in two localities in Colombia suggest quantities between 8.0 and 40.0kg ha⁻¹ of N each month during times without water deficit (Cenicafé, 2007).

The fractions of N available for coffee (NO₃⁻ and NH₄⁺) present considerable variations over time in response to climatic changes and management practices (Ochoa *et al.*, 2003; Sadeghian, 2010), an aspect that leads to the use of OM content as the primary indicator of N levels. In soil analysis calibration research (Figure 2), it was possible to explain relative coffee yield as a function of OM for Colombian conditions (Sadeghian, 2011). The quadratic model suggests that when nitrogen fertilization is suspended for three consecutive years in soils with OM<4.0%, the yield is reduced by 80.0%, and, when the OM>18.0%, the product also decreases because of limitations in mineralization caused by a lower temperature of the sites. Since the highest relative yield value was less than 90.0%, it is possible to interpret that there is always a response to N's supply.

As a process that causes environmental pollution and reduces NUE, N leaching is mainly associated with a nitric fraction (NO₃⁻). Its magnitude increases in more sandy soils and lower CEC and OM contents (Gaines & Gaines, 1994). Hincapié & Henao (2008) and Fenilli *et al.* (2008) argue that under field conditions (Colombia and Brazil, respectively), less than 2.0% of the N applied is lost by leaching; in turn, Babbar & Zak (1995) report losses close to 8.0% in unshaded coffee plantations and 3.0% in plantations under shade for conditions of Costa Rica. Cannavo *et al.* (2013) found higher losses for shaded plantations with *Inga densiflora* Benth. (157.2kg ha⁻¹ year⁻¹ of N), of which 53.0% (83kg ha⁻¹ year⁻¹ of N) corresponded to applied N (250kg ha⁻¹ year⁻¹). In research carried out in Costa Rica, Tully *et al.* (2012) found no differences in N losses due to leaching between organic and conventional management; in contrast, N losses decreased linearly as the biomass of the shade trees increased, a result that suggests the ecosystem service of the trees to




Figure 2. Relationship between soil OM and relative coffee yield (Sadeghian, 2011).

mitigate N leaching. The application of high N rates (400 and 800kg ha⁻¹ year⁻¹), added to the high rainfall, can contribute to more significant leaching of N (up to 13.0% of the applied fertilizer) in irrigated plantations (Bortolotto *et al.*, 2012). It has been shown that there are differences in losses between the soils of the Colombian coffee zone (Figure

3), in particular, due to their clay mineralogy, OM level, and pH rather than soil texture (Arias *et al.*, 2009). To fertilizer sources, more significant leaching has been recorded under laboratory conditions (90.0% of losses as NH₄⁺) when using ammonium sulfate (AS), followed by ammonium nitrate and urea (González & Sadeghian, 2012a) (Figure 4).




Figure 3. Nitrogen losses due to leaching, with and without N (urea) application in four soil-mapping units in Colombia (Arias *et al.*, 2009). Soil mapping unit: Soil mapping units are defined by a group of soils with similar parental material and genetic horizons, with different weathering processes.




Figure 4. Nitrogen losses due to leaching overtime in an Andisol in Colombia due to the application of N sources (González & Sadeghian, 2012a).

The volatilization of $N-NH_3$ is one of the most studied processes in recent years, an aspect that is more related to NUE than air pollution. Although volatilization occurs when applying any source of N, its magnitude is greater with urea. When this source is exposed to the soil surface, it hydrolyzes to form ammonium (NH_4^+) by the action of the enzyme urease. Subsequently, ammonia (NH_3) formation is favored, which is then released into the atmosphere, generating the loss of the element. These losses depend on the following soil properties: pH, CEC, OM, temperature, moisture, and buffer capacity (Dari *et al.*, 2019). Shallow values ($\leq 2.0\%$) are reported for coffee when ammonium sulfate (Fenilli *et al.*, 2008; González & Sadeghian, 2012b) or ammonium nitrate (Dominghetti *et al.*, 2016) are used, and losses between 15.5 and 43.2% with the use of urea (Leal *et al.*, 2007; González & Sadeghian, 2012b; González *et al.*, 2015; Chagas *et al.*, 2016; Dominghetti *et al.*, 2016). The $N-NH_3$ losses are expected to increase with increasing pH, as occurs with the initial urea reaction; the

opposite situation occurs when AS is applied, which tends to generate more acidification due to the nitrification of ammonia. For an Andisol from Colombia, González *et al.* (2015) detected increases of up to 0.5 pH units due to urea application.

Legumes' biological fixation of atmospheric N (N_2) also contributes to N's balance in the system and NUE. The factors that influence the fixation of N_2 are the supply of soil nutrients, pH, legume management, causes that affect photosynthesis, and fertilization with N (Havlin *et al.*, 2017). For Mexican conditions, amounts set by *Inga jinicuil* between <1.0 and $>40.0 \text{ kg ha}^{-1} \text{ year}^{-1}$ are reported (Roskoski, 1982). Legume leaves are generally the primary sink for fixed N (Domenach, 1995); in this regard, $22.7 \text{ kg ha}^{-1} \text{ year}^{-1}$ of N are reported for *Inga densiflora* (Cannavo *et al.*, 2013), and values between 5.0 and 22.0% for Burundi conditions (Snoeck *et al.*, 2000). The cover legumes that grow in the coffee plantations and protect the soil from erosion provide significant amounts of N, of which

part comes from an atmospheric fixation; for example, in the biomass generated by *Arachis pintoï*, there are about 145kg ha⁻¹ (Arango *et al.*, 2009; Rose *et al.*, 2019) and in *Desmodium* spp. 65.0kg ha⁻¹ (Arango *et al.*, 2009). In Brazil, *A. pintoï* in coffee crops was related to an increase in the soil microbial biomass and higher C and N levels, the preceding favored by the low C/N ratio of this coverage, the fast decomposition of its organic residues, and the immobilization of N by microorganisms; *A. pintoï* was also related to the more significant presence of edaphic fauna of the Isoptera order, linked to the high microbial activity in the soil (Lammel *et al.*, 2015). Additionally, Rose *et al.* (2019) argue that the coverage of *A. pintoï* in coffee plantations can offset a part of the external contributions of N while reducing emissions of nitrous oxide (N₂O).

N₂O emitted into the atmosphere through nitrification and denitrification processes is about 300 times more effective than CO₂ in catching heat in the atmosphere; however, the percentage contribution of this gas to global warming, due to the greenhouse effect, is low (IPCC, 2001). N₂O formation occurs under anaerobic conditions, and its magnitude depends on drainage, NO₃⁻ and NO₂⁻ levels, pH, temperature, decomposable OM, quality and quantity of organic residues, nutrient reserves, and texture (Rosenstock *et al.*, 2014; Farquharson, 2016). In most studies on coffee, N₂O values less than 3.0kg ha⁻¹ year⁻¹ are reported (Babbar & Zak, 1995; Fenilli *et al.*, 2008; Ortiz *et al.*, 2018; Montenegro-Ballester, 2019; Rose *et al.*, 2019); nevertheless, in some, up to 12.0kg ha⁻¹ year⁻¹ have been detected (Hergoualc'h *et al.*, 2012; Cannavo *et al.*, 2013; Capa *et al.*, 2015). Variations in N₂O emissions, among other aspects, have been related to

differences in soil N and CO levels (Rose *et al.*, 2019), N rate (Montenegro-Ballester, 2019), source of N (Montenegro, 2020), fertilizer placement (Rose *et al.*, 2019), the water in the soil and rainy seasons (Hergoualc'h *et al.*, 2007; Fenilli *et al.*, 2008; Ortiz *et al.*, 2018; Montenegro-Ballester, 2019). Additionally, more denitrification and N₂O emissions may occur in coffee agroforestry systems than in unshaded plantations (Babbar & Zak, 1995; Hergoualc'h *et al.*, 2012). All this, due to a higher input of N through litterfall and a higher potential mineralization rate of OM in shaded crops. However, the net negative balance of the soil in agroforestry may be lower than in monoculture when the C accumulation in the biomass of the trees is considered (Hergoualc'h *et al.*, 2012).

Climate. The interaction of the elements or components of the climate (temperature, precipitation, air humidity, solar radiation, cloud cover, atmospheric pressure, winds, evaporation, and evapotranspiration) affects the growth of coffee (Jaramillo, 2018) and many of the processes that ultimately govern the NUE (Fageria, 2014; Sarkar & Baishya, 2017). Rain and soil surface temperature (up to 25.0cm deep), which depends on the air temperature (Jaramillo, 2018), are the elements with the most significant influence on the processes that affect N availability. These processes refer to the activity of the microorganisms that mineralize OM, biological fixation, immobilization, nitrification, denitrification, leaching, erosion, volatilization, plant growth, and N absorption and partitioning (Balasubramanian *et al.*, 2004; Weil & Brady, 2017).

The amount and distribution of rainfall show variations between sites and over time (inter and intra-annual), aspects that

must be considered to define fertilization plans, especially regarding application time and the source of N (Sadeghian & Jaramillo, 2016; Sadeghian *et al.*, 2017). Low rainfall is related to limitations in the volume of water containing available N (NO_3^- and NH_4^+) for coffee (Sadeghian *et al.*, 2017), while excessive precipitation increases leaching losses (Sadeghian *et al.*, 2015). The coffee

regions differ considerably in their rainfall even in the same country, as shown in Figure 5 for four localities in the Colombian coffee zone. According to the locality, soil properties, and production system, soil temperature also changes. An example of coffee plantations in full sunlight and with the shade of *Inga* spp. is presented in Figure 6.




Figure 5. Annual variations in rainfall in four municipalities in the Colombian coffee zone. The parentheses denote the department. Original data from Cenicafé (2021).




Figure 6. Soil temperature in coffee plantations unshaded and with the shade of *Inga* spp. in five Colombian soil-mapping units. Higher temperatures are recorded in unshaded coffee plantations than in shady plantations (Cardona & Sadeghian, 2005).

Solar radiation directly affects photosynthesis, which influences the plant's demand for nutrients (Fageria, 1998). In a study developed by Bote *et al.* (2018a), both solar radiation and N supply positively affected some of the attributes of the coffee plant, with positive and negative interactions between the two variables. Bote *et al.* (2018b) also recorded more use of N and higher biomass and production when increasing the rate of N for a higher level of radiation. In this case, N's recovery ranged between 7.0 and 17.0% and decreased with a higher supply of N and increased with the level of radiation. Therefore, it is essential to include the physiological traits of plants in crop breeding programs that improve their ability to tolerate multiple climatic stress factors (Sarkar & Baishya, 2017).

Management. Adopting appropriate soil, plant, climate, and fertilizer management practices improve NUE and favor maximum economic crop yield (Fageria & Baligar, 2005). Most of the environmental components are not controllable; instead, the management seeks to reduce the negative effect of some and take advantage of the benefits of others.

Practices to increase NUE should aim to reduce N losses in the system, improve crops' absorption and use and achieve high productivity (Degaspari *et al.*, 2020). Some of these practices are directly related to the supply of the nutrient through fertilization, specifically, rate and source, time, place and fertilizer placement, and the balance of N with other nutrients; meanwhile, different practices have more indirect effects, for example, productive and N use efficient genotypes, erosion control, liming, use of organic fertilizers, association with legume species, management of pests, diseases and weeds, irrigation, planting densities and spatial arrangements, shade level, planting date and renovation systems. One or more

of the above practices have been cited by Batabyal (2017), Fageria (1998), and Fageria & Baligar (2005).

Nitrogen rate. It is crucial to consider the difference between N management's efficiency and effectiveness. There are different ways to define the NUTUE and NUE; however, they generally resemble that the highest efficiency occurs when the fertilizer is applied at lower rates than the economically optimal rate (Dibb, 2000; Fixen & García, 2006), as illustrated in Figure 7. This last alternative may go against the economic sustainability of coffee farming, and, therefore, it would be necessary to define the priorities in the two management visions.

There are differences between the recommended N rates and the criteria that support this practice. For Colombia, rates between 120 and 300 kg ha⁻¹ year⁻¹ of N are recommended (Sadeghian, 2013), according to the OM content and the production system (affected by planting density and shade level), for Brazil from 50.0 to 450 kg ha⁻¹ year⁻¹, according to the productivity and N foliar concentration (Raj *et al.*, 1996) or based on the crop productivity according to the planting density (Guimarães *et al.*, 1999). In this regard, it is necessary to bear in mind that excess N causes environmental losses and contamination, hence, the care required to select the appropriate rate based on the different soil and plant indicators. The use of SPAD-502 has been evaluated as an indirect measure to estimate foliar N through the chlorophyll reading, to generate indicators that allow adjusting the rates of N and increasing its efficiency. In this regard, while in some cases, the use of this instrument has been successful (Netto *et al.*, 2005; Reis *et al.*, 2009), in others, it has not (Rendón & Sadeghian, 2018), which is why it is currently not widely used, despite its comparative advantages.




Figure 7. Efficiency vs. efficacy for the response of coffee to nitrogen supply. Adapted from Dibb (2000) and Fixen & García (2006).

Nitrogen source. In general, the research results on fertilizer sources do not reveal significant differences in coffee yield, as shown in Figure 8, which is an aspect that is commented on by Malavolta (1993); however, this does not necessarily imply that they cause similar effects on the soil, that their gains and losses are comparable, or that their compatibility in mixtures is the same. To select the appropriate source, it is necessary to jointly consider aspects such as the requirement of the crop, the result of the soil analysis, the interactions of the nutrients and their balance with N, the cost, the residuality, and the availability, among others. Currently, many professionals and companies condemn the use of urea due to its volatilization, frequently ignoring that nitric sources (for example, calcium nitrate, magnesium nitrate, and ammonium nitrate) present more losses due to leaching. Another aspect is the residual acidity, in which urea and ammonium nitrate are comparable. At the same time, AS generates more acidity and can be used more frequently in soils with high pH for coffee (Sadeghian & Duque, 2019).

Application time. In order to increase NUE, it is necessary to apply fertilizers at times of most significant demand for the crop and adequate soil moisture conditions for two or three consecutive months (Malavolta, 1993; Sadeghian & Jaramillo, 2016; Sadeghian *et al.*, 2017). Malavolta (1993) and Favarin *et al.* (2010) are based on the demand for N by the coffee fruit to define the fertilization seasons; in this regard, Sadeghian (2013) draws attention to the issue and clarifies that the condition of soil moisture, governed mainly by rain, prevails over the times of most significant accumulation of N in the fruit if the soil is dry. Furthermore, the requirements of the other organs of the plant in different periods of fruit growth should not be forgotten. In this regard, Fenilli *et al.* (2008) consider that to estimate the efficiency of the recovery of supplied N, it is necessary to take into account the content of N accumulated in other organs of the plant, in addition to the harvested fruits; consequently, 19.1% are reported in the aerial part of the plant, 9.4% in the roots, 23.8% in the litter and 26.3% in the fruits.




Figure 8. Cherry coffee yields in response to three nitrogen sources in an Andisol from Colombia (Sadeghian, 2022). Average values of three rates of each source (100, 200 and 400kg ha⁻¹ year⁻¹). ENTEC® [DMPP (3.4-dimethylpyrazole phosphate, is a nitrification inhibitor)].

Application of nitrogen fertilizers when the soil remains dry can increase soil salinity and cause toxicity in plants, particularly during the establishment stage (Sadeghian *et al.*, 2017). Taking into account the above considerations and based on soil properties, it is generally suggested to supply N and other nutrients two or three times during the year (Cervellini *et al.*, 1986; Martinez *et al.*, 2020; González-Osorio & Sadeghian, 2020); however, when it comes from fertigation, up to 12 applications per year are suggested (Sobreira *et al.*, 2011).

One of the practices to reduce annual fluctuations in coffee production (biennially) has been the fractionation of nitrogen fertilization. Despite the theoretical bases, the results obtained do not reveal any effect (Sanzonowicz *et al.*, 2000).

Nitrogen placement. It is suggested to place the fertilizer where the roots are, bearing in mind both the horizontal and vertical distribution (Malavolta, 1993; Martinez *et al.*,

2020), which varies with plant growth and renewal systems (Sadeghian, 2013). Since the predominant fraction of N in the soil (NO₃⁻) is very mobile, it easily migrates in the profile after its application, so it is suggested to apply it in a dispersed way when fertilizers solids are treated (excludes drip or fluid fertilization).

CONCLUSIONS

The factors that determine NUE are the same that define coffee crop productivity, namely, plant, soil, climate, and management. Despite in recent years, significant advances have been made to search for genetic targets to improve NUE plants, it may be a decade or more before improved NUE coffee varieties are available or used in world coffee farming.

The soil's physical, chemical, and biological properties and climate elements, mainly temperature, precipitation, and solar radiation, affect coffee growth and many of

the processes that ultimately intervene in the functions of the N cycle and affect NUE; nevertheless, some components of these two factors are not fully controllable.

Management of factors that determine NUE should reduce N losses in the system, increase its absorption and use by coffee plants, and maintain or increase yield.

Conflict of interests: The authors declare that there are no conflicts of interest.

BIBLIOGRAPHIC REFERENCES

- Amaral, J. F.; Martinez, H. E.; Laviola, B. G.; Filho, E. I.; Cruz, C. D. (2011a). Eficiência de utilização de nutrientes por cultivares de café. *Ciência Rural*. 41(4): 621-629. doi: <https://doi.org/10.1590/S0103-84782011005000027>
- Amaral, J. F.; Martinez, H. E.; Laviola, B. G.; Tomaz, M. A.; Filho, E. I.; Cruz, C. D. (2011b). Produtividade e eficiência de uso de nutrientes por cultivares de café. *Coffee Science*. 6(1): 65-74.
- Arango, J. G.; Salazar, L. F.; Morales, C. S. (2009). Establecimiento de coberturas vegetales en el cultivo del café. *Suelos Ecuatoriales*. 32(9): 148-151.
- Arias, E.; Sadeghian, S.; Mejía, B.; Morales, C. S. (2009). Lixiviación del nitrógeno en algunos suelos de la zona cafetera y su relación con la textura. *Revista Cenicafé*. 60(3): 239-252.
- Babbar, L. I.; Zak, D. R. (1994). Nitrogen cycling in coffee agroecosystems: net N mineralization and nitrification in the presence and absence of shade trees. *Agriculture, Ecosystems and Environment*. 48: 107-113. doi: [https://doi.org/10.1016/0167-8809\(94\)90081-7](https://doi.org/10.1016/0167-8809(94)90081-7)
- Babbar, L. I.; Zak, D. R. (1995). Nitrogen loss from coffee agroecosystems in Costa Rica: leaching and denitrification in the presence and absence of shade trees. *Journal of Environmental Quality*. 24: 227-233. doi: <https://doi.org/10.2134/jeq1995.00472425002400020003x>
- Balasubramanian, V.; Alves, B.; Aulakh, M.; Bekunda, M.; Cai, Z.; Drinkwater, L.; Mugendi, D.; Kessel, C. V.; Oenema, O. (2004). Crop, environmental, and management factors affecting nitrogen use efficiency. In: Mosier, A. R.; Syers, J. K.; Freney, J. R. (eds.). *Agriculture and the nitrogen cycle: assessing the impacts of fertilizer use on food production and the environment*. pp. 19-33. Washington, USA: Island Press. 344p.
- Batabyal, K. (2017). Nutrient management for improving crop soil, and environmental quality. In: Naeem, M.; Ansari, A. A.; Gill, S. S. (eds.). *Essential plant nutrients: uptake, use efficiency, and management*. pp. 445-464. Cham, Switzerland: Springer. 569p. doi: <https://doi.org/10.1007/978-3-319-58841-4>
- Bess, C. (2020). Developments in Midwestern precision conservation. In: Delgado, J. A.; Gantzer, C. J.; Sassenrath, G. F. (eds.). *Soil and water conservation: a celebration of 75 years*. pp. 286-292. Ankeny, USA: Soil and Water Conservation Society. 331p.
- Bhatla, S. C.; Lal, M. A. (2018). *Plant physiology, development and metabolism*. Singapore: Springer. 1189p. doi: <https://doi.org/10.1007/978-981-13-2023-1>
- Bortolotto, R.; Bruno, I.; Reichardt, K.; Timm, L. C.; Amado, T. J.; Ferreira, A. (2012). Nitrogen fertilizer (¹⁵N) leaching in a central pivot fertigated coffee crop. *Revista Ceres*. 59(4): 466-475. doi: <https://doi.org/10.1590/S0034-737X2012000400006>
- Bote, A. D.; Ayalew, B.; Ocho, F. L.; Anten, N. P.; Vos, J. (2018a). Analysis of coffee (*Coffea arabica* L.) performance in relation to radiation level and rate of nitrogen supply I. Vegetative growth, production and distribution of biomass and radiation use efficiency. *European Journal of Agronomy*. 92: 115-122. doi: <https://doi.org/10.1016/j.eja.2017.10.007>
- Bote, A. D.; Zana, Z.; Ocho, F. L.; Vos, J. (2018b). Analysis of coffee (*Coffea arabica* L.) performance in relation to radiation level and rate of nitrogen supply II. Uptake and distribution of nitrogen, leaf photosynthesis and first bean yields. *European Journal of Agronomy*. 92: 107-114. doi: <https://doi.org/10.1016/j.eja.2017.10.006>

- Bruulsema, T. W.; Witt, C.; García, F.; Li, S.; Rao, N.; Chen, F.; Ivanova, S. (2008). A global framework for fertilizer BMPs. *Better Crops*. 92(2): 13-15.
- Cannavo, P.; Harmand, J. M.; Zeller, B.; Vaast, P.; Ramírez, J. E.; Dambrine, E. (2013). Low nitrogen use efficiency and high nitrate leaching in a highly fertilized *Coffea arabica*-*Inga densiflora* agroforestry system: a ¹⁵N labeled fertilizer study. *Nutrient Cycling in Agroecosystems*. 95: 377-394. doi: <https://doi.org/10.1007/s10705-013-9571-z>
- Capa, D.; Pérez-Esteban, J.; Masaguer, A. (2015). Unsustainability of recommended fertilization rates for coffee monoculture due to high N₂O emissions. *Agronomy for Sustainable Development*. 35: 1551-1559. doi: <https://doi.org/10.1007/s13593-015-0316-z>
- Cardona, D. A.; Sadeghian, S. (2005). Caracterización de la fertilidad del suelo en monocultivos de café (*Coffea Arabica*) y bajo sombrío de guamo (*Inga ssp*). *Colombia Forestal*. 9(18): 87-97. doi: <https://doi.org/10.14483/udistrital.jour.colomb.for.2005.1.a07>
- Cardoso, P. M. (2010). Análise biométrica da eficiência nutricional para nitrogênio em café (*Coffea arabica* L.). Recovered from <http://www.sbicafe.ufv.br:80/handle/123456789/8675>
- Cassman, K. G.; Dobermann, A. R.; Walters, D. T. (2002). Agroecosystems, nitrogen-use efficiency, and nitrogen management. *Ambio*. 31(2): 132-140.
- Cenicafé - Centro Nacional de Investigaciones de Café. (2007). Informe anual Cenicafé 2007. Recuperado de <https://doi.org/10.38141/10783/2007>
- Cenicafé - Centro Nacional de Investigaciones de Café. (2020). Informe Anual Cenicafé 2020. Recovered from <https://doi.org/10.38141/10783/2020>
- Cenicafé - Centro Nacional de Investigaciones de Café. (2021). Registros históricos de lluvia. Recovered from <https://agroclima.cenicafe.org/web/guest/registros-historicos>
- Cervellini, G.; Toledo, S.; Reis, A.; Rocha, T. (1986). Nitrogênio na adubação química do cafeeiro: doses e parcelamentos do nitrocálcio. *Bragantia*. 45(1): 45-55. doi: <https://doi.org/10.1590/S0006-87051986000100006>
- Chagas, W. F.; Guelfi, D.; Caputo, A. L.; Souza, T.; Andrade, A.; Faquin, V. (2016). Ammonia volatilization from blends with stabilized and controlled-released urea in the coffee system. *Ciência e Agrotecnologia*. 40(5): 497-509. doi: <https://doi.org/10.1590/1413-70542016405008916>
- Colodetti, T. V.; Rodrigues, W.; Martins, L.; Tomaz, M. A. (2014). Differential tolerance between genotypes of conilon coffee (*Coffea canephora*) to low availability of nitrogen in the soil. *Australian Journal of Crop Science*. 8(12): 1648-1657.
- Dari, B.; Rogers, C. W.; Walsh, O. S. (2019). Understanding factors controlling ammonia volatilization from fertilizer nitrogen applications. *University of Idaho Extension Bulletin*. 926: 1-4.
- Degaspari, I.; Soares, J.; Montezano, Z. F.; Del Grosso, S.; Vitti, A. C.; Rossetto, R.; Cantarella, H. (2020). Nitrogen sources and application rates affect emissions of N₂O and NH₃ in sugarcane. *Nutrient Cycling in Agroecosystems*. 116: 329-344. doi: <https://doi.org/10.1007/s10705-019-10045-w>
- Dibb, D. W. (2000). The mysteries (myths) of nutrient use efficiency. *Better Crops*. 84(3): 3-5.
- Dobermann, A. R. (2005). Nitrogen use efficiency – state of the art. In: International Fertilizer Industry Association (Compiler). IFA International Workshop on Enhanced-Efficiency Fertilizers. pp. 1-16. Frankfurt, Hesse, Germany: International Fertilizer Industry Association.
- Dobermann, A. R. (2007). Nutrient use efficiency – measurement and management. In: International Fertilizer Industry Association (ed.). Fertilizer best management practices. pp. 1-28. Brussels, Belgium: International Fertilizer Industry Association. 259p.
- Dobermann, A.; Cassman, K. G. (2002). Plant nutrient management for enhanced productivity in intensive grain production systems of the United States and Asia. *Plant and Soil*. 247(1): 153-175. doi: <https://doi.org/10.1023/A:1021197525875>
- Domenach, A. M. (1995). Approche de l'estimation de la fixation symbiotique des arbres par l'utilisation des abondances isotopiques naturelles de l'azote. In: Maillard, P.; Bonhomme, R. (eds.). Utilisation des isotopes stables pour l'étude du fonctionnement des plantes. pp. 159-172. Paris, France: INRA Editions. 360p.

- Dominghetti, A. W.; Guelfi, D.; Guimarães, R. J.; Caputo, A. L.; Spehar, C. R.; Faquin, V. (2016). Nitrogen loss by volatilization of nitrogen fertilizers applied to coffee orchard. *Ciência e Agrotecnologia*. 40(2): 173- 183. doi: <http://dx.doi.org/10.1590/1413-70542016402029615>
- Engels, C.; Kirkby, E.; White, P. (2012). Mineral nutrition, yield and source-sink relationships. In: Marschner, P. (ed.). *Marschner's mineral nutrition of higher plants*. pp. 85-133. Third Edition. United States of America: Academic Press. 651p. doi: <https://doi.org/10.1016/C2009-0-63043-9>
- EUNEP - EU Nitrogen Expert Panel. (2015). Nitrogen use efficiency (NUE) – an indicator for the utilization of nitrogen in agriculture and food systems. Wageningen, Netherlands: EU Nitrogen Expert Panel. 47p.
- Fageria, N. K. (1998). Otimização da eficiência nutricional na produção das culturas. *Revista Brasileira de Engenharia Agrícola e Ambiental*. 2(1): 6-16. doi: <https://doi.org/10.1590/1807-1929/agriambi.v02n01p6-16>
- Fageria, N. K. (2014). Nitrogen management in crop production. Boca Raton, USA: CRC Press. 436p. doi: <https://doi.org/10.1201/b17101>
- Fageria, N. K.; Baligar, V. C. (2005). Enhancing nitrogen use efficiency in crop plants. *Advances in Agronomy*. 88: 97-185. doi: [https://doi.org/10.1016/S0065-2113\(05\)88004-6](https://doi.org/10.1016/S0065-2113(05)88004-6)
- Farquharson, R. (2016). Nitrification rates and associated nitrous oxide emissions from agricultural soils - a synopsis. *Soil Research*. 54(5): 469-480. doi: <https://doi.org/10.1071/SR15304>
- Favarin, J. L.; Tezotto, T.; Neto, A. P.; Pedrosa, A. (2010). Cafeeiro. In: Prochnow, L. I.; Casarin, V.; Stipp, S. R. (eds.) *Boas práticas para uso eficiente de fertilizantes*. pp. 413-467. Volume three. Piracicaba, Brasil: International Plant Nutrition Institute. 467p.
- Fenilli, T. A.; Reichardt, K.; Favarin, J. L.; Bacchi, O.; Silva, A. L.; Timm, L. C. (2008). Fertilizer ¹⁵N balance in a coffee cropping system: a case study in Brazil. *Revista Brasileira de Ciência do Solo*. 32(4): 1459-1469. doi: <https://doi.org/10.1590/S0100-06832008000400010>
- Filho, O. F.; Malavolta, E. (2003). Studies on mineral nutrition of the coffee plant (*Coffea arabica* L. cv. Catuaí Vermelho). LXIV. Remobilization and re-utilization of nitrogen and potassium by normal and deficient plants. *Brazilian Journal of Biology*. 63(3): 481-490. doi: <https://doi.org/10.1590/S1519-69842003000300014>
- Fixen, P. E.; García, F. O. (2006). Effective nutrient management decisions ... looking beyond the next harvest. In: Asociación Argentina de Productores en Siembra Directa (Compiler). *Memorias del Congreso de Asociación Argentina de Productores en Siembra Directa*. pp. 1-11. Rosario, Santa Fe, Argentina: Asociación Argentina de Productores en Siembra Directa.
- Fixen, P. E.; Brentrup, F.; Bruulsema, T.; García, F.; Norton, R.; Zingore, S. (2015). Nutrient/fertilizer use efficiency: measurement, current situation and trends. In: Drechsel, P.; Heffer, P.; Magen, H.; Mikkelsen, R.; Wichelns, D. (eds.) *Managing water and fertilizer for sustainable agricultural intensification*. pp. 8-38. First edition. Paris, France: IFA, IWMI, IPNI, IPI. 257p.
- Gaines, T. P.; Gaines, S. T. (1994). Soil texture effect on nitrate leaching in soil percolates. *Communications Soil Science And Plant Analysis*. 25(13-14): 2561-2570. doi: <https://doi.org/10.1080/00103629409369207>
- González, H.; Sadeghian, S. (2012a). Lixiviación de nitrógeno en suelos de la zona cafetera a partir de diferentes fuentes fertilizantes. *Revista Cenicafé*. 63(1): 111-119.
- González, H.; Sadeghian, S. (2012b). Volatilización del nitrógeno a partir de diferentes fuentes fertilizantes en la etapa de crecimiento vegetativo del café. *Revista Cenicafé*. 63(1): 132-143.
- González, H.; Sadeghian, S.; Medina, R.; Castro, A. (2015). Alternativas para disminuir la volatilización de nitrógeno producida por la fertilización con urea. *Revista Cenicafé*. 66(1): 7-16.
- González-Osorio, H.; Sadeghian, S. (2020). Producción de café en respuesta al fraccionamiento de la fertilización. *Revista Cenicafé*. 71(1): 54-67. doi: <https://doi.org/10.38141/10778/1120>

- Graham, R. D. (1984). Breeding for nutritional characteristics in cereals. In: Tinker, P. B.; Lauchi, A. (eds.) *Advances in plant nutrition*. pp. 57-102. Volume three. New York, USA: Praeger. 320p.
- Guimarães, P. T.; Garcia, A.; Alvarez, V. H.; Prezotti, L. C.; Viana, A.; Miguel, A.; Malavolta, E.; Corrêa, J. C.; Lopes, A.; Nogueira, F.; Monteiro, A.; Oliveira, J. (1999). Cafeeiro. In: Ribeiro, A.; Guimarães, P. T.; Alvarez, V. H. (eds.). *Recomendações para o uso de corretivos e fertilizantes em Minas Gerais*. pp. 289-302. Viosa, Brazil: Comissão de Fertilidade do Solo do Estado de Minas Gerais. 359p.
- Havlin, J. L.; Tisdale, S. L.; Nelson, W. L.; Beaton, J. D. (2017). *Soil fertility and fertilizers: an introduction to nutrient management*. Eighth edition. New Delhi, India: Pearson India. 520p.
- Hawkesford, M.; Horst, W.; Kichey, T.; Lambers, S.; Schjoerring, J.; Møller, I.; White, P. (2012). Function of macronutrients. In: Marschner, P. (ed.). *Marschner's mineral nutrition of higher plants*. pp. 135-189. Third edition. Oxford, UK: Academic Press. 651p. doi: <https://doi.org/10.1016/B978-0-12-384905-2.00006-6>
- Hergoualc'h, K.; Skiba, U.; Harmand, J. M.; Oliver, R. (2007). Processes responsible for the nitrous oxide emission from a Costa Rican Andosol under a coffee agroforestry plantation. *Biology and Fertility of Soils*. 43(6): 787-795. doi: <https://doi.org/10.1007/s00374-007-0168-z>
- Hergoualc'h, K.; Blanchart, E.; Skiba, U.; Catherine, H.; Harmand, J. M. (2012). Changes in carbon stock and greenhouse gas balance in a coffee (*Coffea Arabica* L) monoculture versus an agroforestry system with *Inga densiflora*, in Costa Rica. *Agriculture, Ecosystems and Environment*. 148: 102-110. doi: <https://doi.org/10.1016/j.agee.2011.11.018>
- Hincapié, É.; Henao, M. C. (2008). Efecto de la fertilización del café sobre la composición de la fase líquida en suelos derivados de cenizas volcánicas. Fourteenth edition. Villavicencio, Meta, Colombia: Sociedad Colombiana de la Ciencia del Suelo.
- Hincapié, É.; Salazar, L. F. (2011). Impacto de la erosión sobre las propiedades físicas y químicas del suelo y la producción de café. *Revista Cenicafé*. 62(2): 79-89.
- International Coffee Organization. (2021). Datos históricos de producción total. Recovered from https://www.ico.org/es/new_historical_c.asp
- IPCC - Intergovernmental Panel on Climate Change. (2001). *Climate change 2001: The scientific basis*. NY, USA: Cambridge University Press. 881p.
- Jaramillo, Á. (2018). *El clima de la caficultura en Colombia*. Manizales, Colombia: Centro Nacional de Investigaciones de Café. 206p.
- Lal, R. (2020). Soil erosion and gaseous emissions. *Applied Sciences*. 10(8): 2784. doi: <https://doi.org/10.3390/app10082784>
- Lammel, D. R.; Azevedo, L. C.; Paula, A. M.; Armas, R. D.; Baretta, D.; Cardoso, E. J. (2015). Microbiological and faunal soil attributes of coffee cultivation under different management systems in Brazil. *Brazilian Journal of Biology*. 75(4): 894-905. doi: <http://dx.doi.org/10.1590/1519-6984.02414>
- Lassaletta, L.; Billen, G.; Grizzetti, B.; Anglade, J.; Garnier, J. (2014). 50 year trends in nitrogen use efficiency of world cropping systems: the relationship between yield and nitrogen input to cropland. *Environmental Research Letters*. 9(10): 105011. doi: Recovered from <https://hal.archives-ouvertes.fr/hal-01194829>
- Leal, L. A.; Salamanca-Jiménez, A.; Sadeghian, S. (2007). Pérdidas de nitrógeno por volatilización en cafetales en etapa productiva. *Revista Cenicafé*. 58(3): 216-226.
- Li, H.; Bin, H.; Chu, C. (2017). Nitrogen use efficiency in crops: lessons from *Arabidopsis* and rice. *Journal of Experimental Botany*. 68(10): 2477-2488. doi: <https://doi.org/10.1093/jxb/erx101>
- López, J. C. (2009). Estudio de la eficiencia en el uso del nitrógeno por el café. *Revista Cenicafé*. 60(4): 324-350.
- Machado, L.; Martins, L.; Rodrigues, W.; Ferreira, D.; Côgo, A.; Tomaz, M. A.; Amaral, J. F. (2016). Efficiency and response of conilon coffee genotypes to nitrogen supply. *African Journal of Biotechnology*. 15(35): 1892-1898. doi: <https://doi.org/10.5897/AJB2016.15558>
- Malavolta, E. (1993). *Nutrição mineral e adubação do cafeeiro*. São Paulo, Brazil: Agronômica Ceres. 210p.

- Malavolta, E.; Favarin, J. L.; Malavolta, M.; Cabral, C.; Heinrichs, R.; Silveira, J. S. (2002). Repartição de nutrientes nos ramos, folhas e flores do cafeeiro. *Pesquisa Agropecuária Brasileira*. 37(7): 1017-1022. doi: <https://doi.org/10.1590/S0100-204X2002000700016>
- Mandal, V. K.; Sharma, N.; Raghuram, N. (2018). Molecular targets for improvement of crop nitrogen use efficiency: current and emerging options. In: Shrawat, A.; Zayed, A.; Lightfoot, D. A. (eds.). *Engineering nitrogen utilization in crop plants*. pp. 77-93. Cham, Switzerland: Springer. doi: https://doi.org/10.1007/978-3-319-92958-3_5
- Martinez, H. E.; Souza, B.; Caixeta, E.; Carvalho, F.; Clemente, J. (2020). Water deficit changes nitrate uptake and expression of some nitrogen related genes in coffee-plants (*Coffea arabica* L.). *Scientia Horticulturae*. 267(1): 109254. doi: <https://doi.org/10.1016/j.scienta.2020.109254>
- Martins, L.; Machado, L.; Tomaz, M. A.; Amaral, J. F. (2015). The nutritional efficiency of *Coffea* spp. A review. *African Journal of Biotechnology*. 14(9): 728-734. doi: <https://doi.org/10.5897/AJB2014.14254>
- Masclaux-Daubresse, C.; Daniel-Vedele, F.; Dechorgnat, J.; Chardon, F.; Gaufichon, L.; Suzuki, A. (2010). Nitrogen uptake, assimilation and remobilization in plants: challenges for sustainable and productive agriculture. *Annals of Botany*. 105(7): 1141-1157. doi: <https://doi.org/10.1093/aob/mcq028>
- Mathur, M.; Goel, A. (2017). Quantitative attributes of nutrient uptake and use efficiency. In: Naeem, M.; Ansari, A. A.; Gill, S. S. (eds.). *Essential plant nutrients: uptake, use efficiency, and management*. pp. 81-118. Cham, Switzerland: Springer. 569p. doi: <https://doi.org/10.1007/978-3-319-58841-4>
- Moll, R. H.; Kamprath, E. J.; Jackson, A. (1982). Analysis and interpretation of factors which contribute to efficiency of nitrogen utilization. *Agronomy Journal*. 74(3): 562-564. doi: <https://doi.org/10.2134/agronj1982.0002196200740030037x>
- Montenegro, J. (2020). Efecto de diferentes fuentes de nitrógeno en la emisión de óxido nitroso en plantaciones de café en Costa Rica. *Revista de Ciencias Ambientales*. 54(2): 111-130. doi: <https://doi.org/10.15359/rca.54-2.6>
- Montenegro-Ballester, J. (2019). Respuesta polinómica de la emisión de óxido nitroso en plantaciones de café en Costa Rica. *Revista de Ciencias Ambientales*. 53(2): 1-24. doi: <https://doi.org/10.15359/rca.53-2.1>
- Moody, L.; Bruulsema, T. (2020). From Nutrient Use to Nutrient Stewardship: An Evolution in Sustainable Plant Nutrition. In: Delgado J. A.; Gantzer C. J.; Sassenrath G. F. (eds.). *Soil and water conservation. A Celebration of 75 Years*. pp. 197-202. Ankeny, Iowa: Soil and Water Conservation Society. 332p.
- Netto, A.; Campostrini, E.; Oliveira, J.; Bressan-Smith, R. E. (2005). Photosynthetic pigments, nitrogen, chlorophyll a fluorescence and SPAD-502 readings in coffee leaves. *Scientia Horticulturae*. 104(2): 199 - 209. doi: <https://doi.org/10.1016/j.scienta.2004.08.013>
- Ochoa, W.A.; Suarez V. S.; Sadeghian K. S. (2003). Variabilidad espacial del Nitrógeno disponible en andisoles de la zona cafetera colombiana. *Revista Cenicafé*. 54(2):179-189.
- Ortiz, G. D.; Neegaard, A.; Vaast, P.; Suárez-Villanueva, V.; Oelofse, M.; Rosenstock, T.S. (2018). Multi-scale measurements show limited soil greenhouse gas emissions in Kenyan smallholder coffee-dairy systems. *Science of the Total Environment*. 626(1): 328-339. doi: <https://doi.org/10.1016/j.scitotenv.2017.12.247>
- Pilbeam, D. J. (2015). Nitrogen. In: Barker, A. V.; Pilbeam, D. J. (eds.). *Handbook of plant nutrition*. pp. 17-63. 2 Ed. Boca Ratón, Florida: CRC press. 773p.
- Prezotti, L. C.; Bragança, S. M. (2013). Acúmulo de massa seca, N, P, e K em diferentes materiais genéticos de café conilon. *Coffee Science*. 8(3): 284-294.
- Prezotti, L. C.; Novais, R.; Alvarez, V. H.; Cantarutti, R. B.; Barros, N. F. (2000). Adubação de formação e manutenção de cafezais (sistema para recomendação de fertilizantes e corretivos de solo para a cultura do café arabica). In: Zambolim, L. (ed.). *Café: Produtividade, qualidade e sustentabilidade*. pp. 125-147. Viçosa, Brazil: Universidade Federal de Viçosa. 395p.

- Raij, B. (2011). Fertilidade do solo e manejo de nutrientes. Piracicaba: International Plant Nutrition Institute. 420p.
- Raij, B.; Cantarella, H.; Quaggio, J. A.; Furlani, Â. M. (1996). Boletim Técnico N° 100: recomendações de adubação e calagem para o estado de São Paulo. Second edition. Campinas, Brazil: Instituto Agronômico-FUNDAG. 285p.
- Reis, A.; Favarin, J. L.; Malavolta, E.; Júnior, J. L.; Moraes, M. (2009). Photosynthesis, chlorophylls, and SPAD readings in coffee leaves in relation to nitrogen supply. *Communications in Soil Science and Plant Analysis*. 40(9-10): 1512-1528. doi: <https://doi.org/10.1080/00103620902820373>
- Rendón, J. R.; Sadeghian, S. (2018). Aplicación de índices espectrales para identificar necesidades de fertilización nitrogenada en café. *Revista Cenicafé*. 69(1): 7-15.
- Rose, T. J.; Kearney, L. J.; Morris, S.; Zwieten, L. V.; Erler, D. V. (2019). Pinto peanut cover crop nitrogen contributions and potential to mitigate nitrous oxide emissions in subtropical coffee plantations. *Science of the total Environment*. 656: 108-117. doi: <https://doi.org/10.1016/j.scitotenv.2018.11.291>
- Rosenstock, T. S.; Tully, K. L.; Arias-Navarro, C.; Neufeldt, H.; Butterbach-Bahl, K.; Verchot, L. V. (2014). Agroforestry with N₂-fixing trees: sustainable development's friend or foe?. *Current Opinion in Environmental Sustainability*. 6: 15-21. doi: <https://doi.org/10.1016/j.cosust.2013.09.001>
- Roskoski, J. P. (1982) Nitrogen fixation in a Mexican coffee plantation. *Plant and Soil*. 67: 283-291. doi: <https://doi.org/10.1007/BF02182775>
- Sadeghian, S. (2008). Fertilidad del suelo y nutrición del café en Colombia: guía práctica. *Boletín Técnico Cenicafé*. 32: 1-43.
- Sadeghian, S. (2009). Calibración de análisis de suelo para N P K y Mg en cafetales al sol y bajo semisombra. *Revista Cenicafé*. 60(1): 7-24.
- Sadeghian, S. (2010). Evaluación de la fertilidad del suelo para una adecuada nutrición de los cultivos. Caso café. *Suelos Ecuatoriales*. 41(1): 46-64.
- Sadeghian, S. (2011). Respuesta de cafetales al sol y bajo semisombra a nitrógeno y su relación con la materia orgánica del suelo. *Revista Facultad Nacional de Agronomía Medellín*. 64(1): 5781-5791.
- Sadeghian, S. (2013). Nutrición de cafetales. In: Centro Nacional de Investigaciones de Café (eds.). *Manual del cafetero colombiano: investigación y tecnología para la sostenibilidad de la caficultura*. pp. 85-116. Volume 2. Manizales. Colombia: Centro Nacional de Investigaciones de Café. 352p.
- Sadeghian, S. (2014). Manejo integrado de nutrientes para una caficultura sostenible. *Suelos Ecuatoriales*. 44(2): 74-89.
- Sadeghian, S.; González, H.; Arias, E. (2015). Lixiviación de nutrientes en suelos de la zona cafetera: Prácticas que ayudan a reducirla. *Boletín Técnico Cenicafé*. 40: 1-34.
- Sadeghian, S.; Jaramillo, Á. (2016). Nutrición de los cafetales en Colombia, en escenarios de La Niña. *Avances Técnicos Cenicafé*. 473: 1-12.
- Sadeghian, S.; Jaramillo, Á.; Duque, H. (2017). Nutrición de los cafetales en Colombia en escenarios El Niño. *Avances Técnicos Cenicafé*. 477: 1-12.
- Sadeghian, S.; Duque, H. (2019). Fertilizantes minerales para café en Colombia: consideraciones técnicas. *Avances Técnicos Cenicafé*. 503: 1-8. doi: <https://doi.org/10.38141/10779/0503>
- Sadeghian, S.; Díaz-Marín, C. (2020). Corrección de la acidez del suelo: alteraciones químicas del suelo. *Revista Cenicafé*. 71(1): 7-20. doi: <https://doi.org/10.38141/10778/1116>
- Sadeghian, S. (2022). Nutrición del café. Consideraciones para el manejo de la fertilidad del suelo. Bogota: Cenicafé. <https://doi.org/10.38141/cenbook-0017>
- Sanzonowicz, C.; Sampaio, J. B.; Nazareno, R. (2000). Efeito de dose e parcelamento do nitrogênio na flutuação bienal da produção do cafeeiro no cerrado. Recovered from <http://www.sbicafe.ufv.br/handle/123456789/756>
- Sarkar, D.; Baishya, L. K. (2017). Nutrient use efficiency. In: Naeem, M.; Ansari, A. A.; Gill, S. S.

- (eds.). Essential plant nutrients: uptake, use efficiency, and management. pp. 119-146. Cham, Switzerland: Springer. 569p. doi: <https://doi.org/10.1007/978-3-319-58841-4>
- Sinha, V.; Jangam, A.; Raghuram, N. (2020). Biological determinants of crop nitrogen use efficiency and biotechnological avenues for improvement. In: Sutton, M. A.; Mason, K. E.; Bleeker, A.; Hicks, W. K.; Masso, C.; Raghuram, N.; Reis, S.; Bekunda, M. (eds.). Just Enough Nitrogen: perspectives on how to get there for regions with too much and too little nitrogen. pp. 157-171. Cham, Switzerland: Springer. 608p. doi: <https://doi.org/10.1007/978-3-030-58065-0>
- Snoeck, D.; Zapata, F.; Domenach, A. M. (2000). Isotopic evidence of the transfer of nitrogen fixed by legumes to coffee trees. *Biotechnology, Agronomy, Society and Environment*. 4(2): 95-100.
- Sobreira, F.; Guimarães, R. J.; Colombo, A.; Scalco, M. S.; Carvalho, J. (2011). Adubação nitrogenada e potássica de cafeeiro fertirrigado na fase de formação, em plantio adensado. *Pesquisa Agropecuária Brasileira*. 46(1): 9-16. doi: <https://doi.org/10.1590/S0100-204X2011000100002>
- Soil Science Society of America. (2008). Glossary of soil science terms 2008. Madison, USA: Soil Science Society of America. 84p. doi: <https://doi.org/10.2136/2008.glossarysoilscienceterms>
- Taiz, L.; Zeiger, E.; Møller, I. M.; Murphy, A. (2021). *Fundamentos de Fisiologia Vegetal*. 6 ed. Porto Alegre, Brasil: Artmed Editora. 584p.
- Tully, K. L.; Lawrence, D.; Scanlon, T. M. (2012). More trees less loss: Nitrogen leaching losses decrease with increasing biomass in coffee agroforests. *Agriculture, Ecosystems and Environment*. 161: 137-144. doi: <https://doi.org/10.1016/j.agee.2012.08.002>
- Weil, R. R.; Brady, N. C. (2017). *The nature and properties of soils*. Fifteenth global edition. Harlow, England: Pearson. 1104p.
- World Meteorological Organization. (2019). WMO greenhouse gas bulletin #15: the state of greenhouse gases in the atmosphere based on global observations through 2018. Recovered from <https://reliefweb.int/report/world/wmo-greenhouse-gas-bulletin-state-greenhouse-gases-atmosphere-based-global-0>