

Antimicrobial and molecular resistance profiles of bacterial isolates from poultry production environment

Perfiles de resistencia antimicrobiana y molecular de aislados bacterianos del entorno de producción avícola.

Marina Mota Fernandes¹, Endria Virgínia Alves Silva², Beatriz Barbosa Reis³, Gessyk Monteiro Marques⁴, Tullio Teixeira Deusdará⁵, Hélio de Sousa Brito⁶, Marcela Oliveira Ferreira⁷, Keytte Fernanda Vieira Silva⁸, Ana Angelita Sampaio Baptista⁹, Vladimir Pinheiro do Nascimento¹⁰, Alex Sander Rodrigues Cangussu¹¹

Authors Data

1. Student, M.Sc., Universidade Federal do Tocantins, Gurupi, Brasil, fmarinamota@gmail.com, <https://orcid.org/0000-0002-7937-590X>
2. Student, Universidade Federal do Tocantins, Gurupi, Brasil, endriavirginia@mail.uft.edu.br, <https://orcid.org/0009-0000-1011-1095>
3. Student, Universidade Federal do Tocantins, Gurupi, Brasil, reis.beatriz@mail.uft.edu.br, <https://orcid.org/0009-0006-6864-1420>
4. Student, Universidade Federal do Tocantins, Gurupi, Brasil, gessykmonteiro2203@gmail.com, <https://orcid.org/0009-0003-8174-2320>
5. Researcher, Ph.D., Universidade Federal do Tocantins, Gurupi, Brasil, tullio@uft.edu.br, <https://orcid.org/0000-0002-7566-0631> (Corresponding Author)
6. Researcher, M.Sc., Universidade Federal do Tocantins, Gurupi, Brasil, hellyosousa1245@gmail.com, <https://orcid.org/0000-0002-6785-0267>
7. Student, Universidade Federal do Tocantins, Gurupi, Brasil, ferreira.marcela@mail.uft.edu.br, <https://orcid.org/0009-0002-1603-0776>
8. Researcher, Universidade Federal do Tocantins, Gurupi, Brasil, keyttefernanda@hotmail.com, <https://orcid.org/0009-0007-5434-5169>
9. Professor. Dr., Universidade Estadual de Londrina, Londrina, Brasil, anaangelita@uel.br, <https://orcid.org/0000-0003-0699-7246>
10. Professor. Ph.D., Universidade Federal do Rio Grande do Sul, Porto Alegre, Brasil, vladimir@ufrgs.br, <https://orcid.org/0000-0002-7720-3274>
11. Professor. Ph.D., Universidade Federal do Tocantins, Gurupi, Brasil, alexscangussu@mail.uft.edu.br, <https://orcid.org/0000-0003-2664-8027>



Cite: Fernandes, M.M.; Silva, E.V.A.; Reis, B.B.; Marques, G.M.; Deusdará, T.T.; Brito, H.S.; Ferreira, M.O.; Silva, K.F.V.; Baptista, A.A.S.; Nascimento, V.P.; Cangussu, A.S.R. (2024). Antimicrobial and molecular resistance profile of bacterial isolates from poultry production environment. *Revista de Ciencias Agrícolas*. 41(3): e3240.

<http://doi.org/10.22267/rcia.20244103.240>

Received: June 18 2024

Accepted: November 15 2024

ABSTRACT

Poultry farming is a relevant industry in Brazil, providing essential animal protein. However, such commercial activity faces significant challenges from bacterial-based diseases, which can impact both animal health and quality product. In this context, this study aimed to identify bacterial strains and evaluate their antimicrobial resistance profiles in poultry farms in southern Tocantins, Brazil. Samples were collected from different stages of poultry production, and the antimicrobial

susceptibilities were assessed through antibiogram tests, where bacterial isolates were exposed to 17 antibiotics commonly used in bacterial infection treatment. Bacterial isolates were identified by 16S rRNA gene sequencing, and phylogenetic analysis was used to group them based on genetic similarities. Five isolates were identified: *Glutamicibacter creatinolyticus* (IGA1), *Enterococcus gallinarum* (IGA2), *Enterobacter mori* (IGA3), *Lysinibacillus fusiformis* (IGA4), and *Enterococcus faecalis* (IGA5). Antibiotic susceptibility tests revealed significant variations in resistance profiles, with some isolates exhibiting multidrug-resistant (MDR) phenotypes. IGA1 was classified as multidrug-resistant, showing resistance to imipenem, meropenem, ceftazidime, trimethoprim, and doxycycline. IGA2 exhibited resistance to β -lactam antibiotics, including ceftazidime, cefepime, doxycycline, imipenem, meropenem, piperacillin-tazobactam, and tetracycline. IGA3 was resistant to aztreonam and trimethoprim, while IGA4 showed resistance to ceftazidime and cefepime but sensitivity to ciprofloxacin and linezolid. IGA5 did not show resistance to any of the tested antimicrobials. Collectively, our findings highlight bacterial diversity and antibiotic resistance in poultry farms, reinforcing the need for continuous monitoring and effective antimicrobial control strategies to promote animal health, ensure food safety, and prevent the transmission of multidrug-resistant bacteria to humans.

Keywords: Animal health; food safety; multidrug-resistant; poultry farming; strains; 16S rRNA.

RESUMEN

La avicultura es una industria relevante en Brasil, porque proporciona una fuente esencial de proteína animal. Sin embargo, esta enfrenta desafíos significativos debido a enfermedades bacterianas que pueden afectar la salud animal y la calidad del producto. En este contexto, este estudio tuvo como objetivo identificar cepas bacterianas y evaluar sus perfiles de resistencia a los antimicrobianos en granjas avícolas del sur de Tocantins, Brasil. Se recolectaron muestras en diferentes etapas de la producción avícola, y la susceptibilidad antimicrobiana se evaluó mediante pruebas de antibiograma, exponiendo aislados bacterianos a 17 antibióticos comúnmente utilizados en el tratamiento de infecciones bacterianas. Los aislados fueron identificados mediante la secuenciación del gen 16S rRNA, y se empleó análisis filogenético para agruparlos según similitudes genéticas. Se identificaron cinco aislados: *Glutamicibacter creatinolyticus* (IGA1), *Enterococcus gallinarum* (IGA2), *Enterobacter mori* (IGA3), *Lysinibacillus fusiformis* (IGA4) y *Enterococcus faecalis* (IGA5). Las pruebas de susceptibilidad revelaron variaciones significativas en los perfiles de resistencia, con algunos aislados presentando fenotipos multirresistentes (MDR). IGA1 fue clasificado como multirresistente, mostrando resistencia a imipenem, meropenem, ceftazidima, trimetoprima y doxiciclina. IGA2 mostró resistencia a antibióticos β -lactámicos, incluyendo ceftazidima, cefepima, doxiciclina, imipenem, meropenem, piperacilina-tazobactam y tetraciclina. IGA3 fue resistente a aztreonam y trimetoprima, mientras que IGA4 mostró resistencia a ceftazidima y cefepima, pero sensibilidad a ciprofloxacino y linezolid. IGA5 no mostró resistencia a ninguno de los antimicrobianos probados. Colectivamente, nuestros hallazgos destacan la diversidad bacteriana y perfiles de resistencia a los antibióticos en las granjas avícolas, reforzando la necesidad de monitoreo continuo y estrategias efectivas de control antimicrobiano para promover la salud animal, garantizar la seguridad alimentaria y prevenir la transmisión de bacterias multirresistentes a seres humanos.

Palabras clave: Avicultura; Cepas bacterianas; resistencia a antimicrobianos; sanidad animal; seguridad alimentaria; 16S rRNA.

INTRODUCTION

The poultry protein industry in Brazil is of significant global importance, ranking as the second-largest chicken meat producer in the world. Recent data reveals that the annual production of chicken meat has reached 14.524 million tons, which is exported to 151 different nations, resulting in an export value of \$7.6 billion (ABPA, 2022).

Due to the high demand in this sector, the poultry industry relies on raising large numbers of birds. In this context, it is also necessary to adopt effective alternatives for disease prevention and control and improve the growth performance of poultry. Bacterial infections are among the most common diseases in poultry farming, capable of impacting animal health and product quality, in addition to posing a risk to public health. Examples of bacterial poultry diseases include fowl typhoid (*Salmonella gallinarum*), colibacillosis (*Escherichia coli*), pasteurellosis (*Pasteurella multocida*), infectious coryza (*Avibacterium paragallinarum*), and others (EMBRAPA, 2021). These diseases also raise substantial concerns as they have the potential to cause significant economic losses, affecting both producers and the economy as a whole (Gutarowska *et al.*, 2018; Paudel *et al.*, 2021; Gržinić *et al.*, 2023).

Strategies aimed at reducing the incidence of diseases include the prophylactic use of antimicrobials in humans and animals to combat bacterial infections. Additionally, these agents are administered sub-therapeutically as feed additives, aiming to promote animal growth. Throughout the twentieth century, the use of antibiotics was a significant factor in reducing mortality rates associated with infectious diseases. However, the indiscriminate and inappropriate use of these drugs emerged as one of the main underlying causes of the emergence of multidrug-resistant (MDR) bacteria (Moyane *et al.*, 2013; Baynes *et al.*, 2016; Manyi-Loh *et al.*, 2018; Berghiche *et al.*, 2018; Arbab *et al.*, 2021; Moretto *et al.*, 2021).

The molecular profile and antimicrobial resistance of bacterial isolates are essential for understanding the genetic diversity and resistance of microorganisms (Arbab *et al.*, 2021; Deusdará *et al.*, 2023). Molecular identification involves techniques such as PCR and whole-genome sequencing, which are based on DNA amplification. PCR is used to amplify specific regions of each species, employing different primers, including genes encoding the 16S rRNA and 23S rRNA subunits (Zarrilli *et al.*, 2013). This approach is crucial for identifying pathogens, assessing treatment efficacy, and implementing preventive measures (Gomes *et al.*, 2019; Arbab *et al.*, 2021). Additionally, it helps reduce the risk of the spread of multidrug-resistant pathogens, benefiting animal health, food safety, and public health (Fardsanei *et al.*, 2017; Moretto *et al.*, 2021).

Given the challenges associated with poultry farming, producers need to develop and implement comprehensive prevention plans. This study aims to identify pathogenic microorganisms and analyze the antibiotic susceptibility of strains isolated from poultry

environments in the southern region of Tocantins, Brazil. Consequently, this approach makes a significant contribution to promoting animal health and developing appropriate public health strategies. Furthermore, the study fosters knowledge dissemination, as data on the evaluation of antibiotic-resistant bacteria in poultry farms in southern Tocantins is scarce in the literature.

MATERIALS AND METHODS

Sample Collection and Processing. The samples were collected from chicken poultry farms of the species *Gallus gallus domesticus* in rural property Tocantins, Brazil (11°47'48" S and 49°31'44" W). These were identified with sequential codes of poultry farm isolates: IGA1, IGA2, IGA3, IGA4, and IGA5. These were obtained using sterile microbiological swabs at different locations, such as soil, chicken coop floor, and fecal samples, which were subsequently stored in sterile containers and transported to the laboratory. Sample processing was carried out immediately, and the samples were cultured in a Brain Heart Infusion (BHI) culture medium at 37°C for 24 hours in a Biochemical Oxygen Demand (BOD) incubator. After incubation, the resulting colonies were identified using the 16S RNA gene and the analysis of the antibiotic susceptibility profile.

DNA Extraction, Amplification, and Sequencing. Genomic DNA extraction from bacterial isolates was performed using a commercial kit (Promega, Madison, WI, USA) and silica columns. In summary, DNA was extracted from the collected isolates and added to sterile tubes. The biological material was centrifuged at 14,000 rpm for 1 minute, and the supernatant was removed. The material was homogenized in a lysis solution with extraction buffer and proteinase K, incubated at 55 °C for 30 minutes, and centrifuged at 12,000 rpm for 3 minutes. The supernatant was transferred to a new tube containing isopropanol, and after centrifugation at 12,000 rpm for 2 minutes, the precipitated DNA was collected. The quantity and quality of the extracted DNA were assessed by agarose gel electrophoresis (1%) with a molecular weight marker and intercalating dye. DNA purification involved cell lysis, DNA binding to the silica columns of the kit, washing to remove impurities, and elution of the purified DNA in the elution buffer. The obtained DNA was stored at -20 °C for later analysis.

Amplification of the 16S rRNA gene was carried out using specific primers that recognize conserved sequences at the ends of the gene. The reaction mix contained 10 µL of reaction buffer, 2 µL of universal primers 27F (5'-AGAGTTTGATCMTGGCTCAG-3') and 1492R (5'-GGTACCTTGTACGACTT-3') (10 µM each), 1 µL of dNTPs (10 mM), 0.5 µL of Taq DNA polymerase (5 U/µL), and 1 µL of genomic DNA. The mix was pipetted into PCR tubes and placed in a thermal cycler for amplification.

The amplified DNA was subjected to a sequencing reaction using fluorescent chain terminators (Sanger *et al.*, 1977). The sequencing reaction was prepared with 10 µL of amplified DNA sample, 4 µL of sequencing buffer, 2 µL of sequencing primer (10 µM), and

4µL of BigDye Terminator v3.1 kit from Thermo Fisher Scientific. The mix underwent a thermal cycle: 1 minute at 96°C, followed by 25 cycles of 10 seconds at 96°C, 5 seconds at 50°C, and 4 minutes at 60°C. After sequencing, the mixture was purified using the BigDye XTerminator kit (Thermo Fisher Scientific), precipitated with ethanol, washed with 70% ethanol solution, resuspended in ultrapure water, and subjected to capillary electrophoresis in the Applied Biosystems 3730XL sequencer (Thermo Fisher Scientific). The obtained sequences were analyzed with Sequencing Analysis v5.4 software, with quality assessment using Phred software. Low-quality sequences (Phred < 20) were removed from the analysis, and the remaining sequences were aligned and assembled into contigs using CodonCode Aligner v.11.0.3 software.

Characterization of the antibiotic susceptibility profile. The disk diffusion method was employed to evaluate bacterial susceptibility and resistance classification to antibiotics, following the guidelines of the Clinical and Laboratory Standards Institute (CLSI, 2020). The isolates were cultured in 100 mL of TSB broth (Soybean Tryptone Broth) under agitation at 120 rpm at 37°C for 8 hours. Subsequently, the growth was adjusted to 10⁸ CFU/mL and seeded on Mueller-Hinton agar (Kasvi, Italy). A disk of antibiotics was then inoculated onto the plate and incubated at 37°C for 18 to 24 hours to record the formation of the inhibition zone. The antibiotics (Sensifar®-Brazil) tested were: ampicillin/sulbactam (20µg), amikacin (30µg), ceftazidime (30µg), cefepime (30µg), ciprofloxacin (5µg), cotrimoxazole (25µg), gentamicin (10µg), imipenem (10µg), levofloxacin (5µg), linezolid (30µg), meropenem (10µg), minocycline (30µg), norfloxacin (10µg), piperacillin/tazobactam (110µg), polymyxin B (330 IU), tetracycline (30µg), vancomycin (30µg). These antibiotics allow for a broader assessment of pathogen resistance, providing a better understanding of the effectiveness of treatments. The inhibition zone was measured manually using a metric ruler across the zone inhibition around each antibiotic and classified in resistance to ≥ 1 antimicrobial agent and in ≥ 3 antimicrobial categories considered MDR (Multidrug Resistance) (Magiorakos *et al.*, 2012; Deusdará *et al.*, 2023).

Bioinformatics analysis. The DNA sequences are imported into CodonCode Aligner V4.2.7 software (LI-COR®) and undergo a series of steps, including the identification of overlapping regions, the removal of low-quality regions, and the clustering of similar sequences to produce a consensus sequence. To identify and compare DNA sequences in the NCBI and RDP databases, sequences were first uploaded in FASTA or GenBank format using search tools such as BLAST (Basic Local Alignment Search Tool) and the Ribosomal Database Project (RDP) (Altschul *et al.*, 1990; Cole *et al.*, 2014).

The construction of the phylogenetic tree by dendrogram was performed using MEGA 11 software using multiple alignments of DNA sequences with the MUSCLE algorithm and the analysis of evolution models with the Bayesian information criterion (BIC). The phylogenetic tree was constructed using the Neighbor-Joining (NJ) method using the distance matrix corrected by the chosen model, and finally, the robustness of the tree was assessed through bootstrap analysis with 1000 replicates. The resulting tree was visualized and edited in FigTree software (Kumar *et al.*, 2018).

RESULTS AND DISCUSSION

Bacterial identification using 16S rRNA. A total of five isolates were identified from biological samples (feces) and environmental samples (floor and perch) of poultry farms (Table 1). Isolates IGA1, IGA2, IGA3, IGA4, and IGA5 were identified through the sequencing of the 16S rRNA gene and were found to belong to different bacterial genera (Table 1). In the phylogenetic analysis, isolate IGA1 exhibited 18 characteristic nucleotide sequences of the species *G. creatinolyticus* (Table 1; Figure 1). A total of 1.408 positions in the final dataset of isolate IGA1 showed 99.4% similarity to *G. creatinolyticus* in BLAST, clustering in the clade with 100% bootstrap support (Figure 1). The isolate IGA2 presented 15 characteristic nucleotide sequences (Table 1; Figure 2). In the final dataset, 1.437 positions of the isolate showed 99.86% similarity to *E. gallinarum* based on BLAST analysis, clustering within a clade that was supported by 88% bootstrap (Figure 2). IGA3 exhibited 18 distinctive nucleotide sequences characteristic of the genus *Enterobacter* sp. There were a total of 1414 positions in the final dataset of isolate IGA3 that showed 99.5% similarity to *Enterobacter mori* species in BLAST, clustering in the clade with 86% bootstrap support (Table 1; Figure 3). IGA4 displayed 18 characteristic nucleotide sequences of the species *L. fusiformis* (Table 1; Figure 4). There were a total of 1422 positions in the final dataset of isolate IGA4, demonstrated 99.72% similarity to the bacterium *L. fusiformis* in BLAST, clustering in the clade with 91% bootstrap support (Figure 4). IGA5 showed 21 characteristic nucleotide sequences of the genus and species *E. faecalis*. The final dataset, comprising 1.427 positions, showed 100% similarity (Table 1; Figure 5).

Antibiotic susceptibility testing. The susceptibility and resistance profiles of the five isolates revealed varied patterns for the different antimicrobial agents tested. Isolate IGA1 demonstrated resistance to the antimicrobials imipenem (IPM10), meropenem (MER10), ceftazidime (CAZ30), trimethoprim (TRI5), and doxycycline (DOX30) from the cephalosporin, carbapenem, sulfonamide, and tetracycline classes, thus being classified as MDR (multidrug-resistant) (Table 1). IGA2 showed resistance to different antimicrobial agents tested, including ceftazidime (CAZ30), cefepime (CPM30), doxycycline (DOX30), imipenem (IPM10), meropenem (IPM10), piperacillin-tazobactam (PPT110), and tetracycline (TET30), which belong to the cephalosporin, carbapenem, penicillin, and tetracycline classes (Table 1). However, it is important to note that isolate IGA2 exhibited resistance to β -lactam antibiotics (cephalosporin, carbapenem, and penicillin). IGA3 showed resistance to the antimicrobials aztreonam (ATM30) and trimethoprim (TRI5), while no resistance was observed for the others (Table 1). Isolate IGA4 demonstrated resistance to only two antimicrobials, ceftazidime (CAZ30) and cefepime (CPM30), from the cephalosporin class, and sensitivity to most antimicrobials, including those belonging to the fluoroquinolone class such as ciprofloxacin (CIP5) and norfloxacin (NOR10). Additionally, it showed sensitivity to linezolid (LNZ30) and gentamicin (GEN10). On the other hand, strain IGA5 did not exhibit resistance to any tested antimicrobials (Table 1).

Table 1. Identification of microorganisms by 16S rRNA sequencing compared to GenBank NCBI and antimicrobial resistance.

Isolate (Codes)	Scientific Name	Number GenBank accession	Query cover (%)	Identity (%)	Number GenBank accession 16S rRNA gene partial sequence	Antimicrobial resistance	*MDR
IGA1	<i>G. creatinolyticus</i>	NR_036769.1	98	99.4	OR625203	IPM, MER, CAZ, TRI, DOX	P
IGA2	<i>E. gallinarum</i>	NR_104559.2	100	99.86	OR625204	IPM, MER, CAZ, CPM, DOX, PPT, TET	N
IGA3	<i>Enterobacter</i> sp.	NR_116430.1	100	99.51	OR625205	ATM, TRI	N
IGA4	<i>L. fusiformis</i>	NR_042072.1	100	99.72	OR625206	CAZ, CPM	N
IGA5	<i>E. faecalis</i>	NR_115765.1	99	100	OR625202	-	N

*Antibiotics: imipenem (IPM), meropenem (MER); ceftazidime (CAZ), cefepime (CPM), trimethoprim (TRI), doxycycline (DOX), aztreonam (ATM), piperacillin/tazobactam (PPT), tetracycline (TET). *Multidrug-resistant bacteria. P: Positive; N: Negative.


**Figure 1.** Phylogenetic trees obtained by the neighbor-joining method with the Jukes-Cantor parameter and Bootstrap of 1000 replicas of 16S rRNA gene sequences of the isolate 2023.0290.16S (IGA1) with GenBank accession number OR625203.



Figure 2. Phylogenetic trees obtained by the neighbor-joining method with the Jukes-Cantor parameter and Bootstrap of 1000 replicas of 16S rRNA gene sequences of the isolate 2023.0291.16S (IGA2) with GenBank accession number OR625204.




Figure 3. Phylogenetic trees obtained by the neighbor-joining method with the Jukes-Cantor parameter and Bootstrap of 1000 replicas of 16S rRNA gene sequences of the isolate 2023.0292.16S (IGA3) with GenBank accession number OR625205.




Figure 4. Phylogenetic trees obtained by the neighbor-joining method with the Jukes-Cantor parameter and Bootstrap of 1000 replicas of 16S rRNA gene sequences of isolate 2023.0293.16S (IGA4) with GenBank accession number OR625206.




Figure 5. Phylogenetic trees obtained by the neighbor-joining method with the Jukes-Cantor parameter and Bootstrap of 1000 replicas of 16S rRNA gene sequences of isolate 2023.0289.16S (IGA4) with GenBank accession number OR625206.

Bacterial infections can significantly impair animal health and performance, leading to serious complications, especially when there is resistance to multiple antimicrobials (MDR), which complicates treatment and increases the risks of more severe outbreaks. Therefore, evaluating the resistance profile of bacteria in poultry farms is important since they can be sources of pathogenic microorganisms (Lateef *et al.*, 2015; Deusdará *et al.*, 2023). Thus, using the 16S rRNA gene sequencing, it is possible to evaluate the presence of species of prokaryotic organisms, such as bacteria, fungi, and archaea, and its ability to provide valuable phylogenetic and taxonomic information about microbial diversity, a relevant tool for identifying pathogens in poultry farms (Deusdará *et al.*, 2023).

In the scope of this study, these approaches were implemented, allowing for the characterization of antibiotic resistance. Furthermore, accurate identification of the investigated microorganisms validates the reliability of the 16S rRNA gene sequencing technique for constructing the phylogenetic profile and determining the isolates *G. creatinolyticus* (IGA1), *E. gallinarum* (IGA2), *Enterobacter sp.* (IGA3), *L. fusiformis* (IGA4), and *E. faecalis* (IGA5). *G. creatinolyticus* (IGA1) exhibited multidrug-resistant phenotypes to antibiotics. Is considered ubiquitous, as it can be found in various habitats, including different types of soil, clinical samples, cheeses, and plants, and plays an important role in various biotechnological processes. However, when pathogenic, it is associated with urinary tract infections and bacteremia. This species may be related to opportunistic infections in humans and animals, especially in immunocompromised individuals, and its presence in infections may have significant clinical implications (Hou *et al.*, 1998; Santos *et al.*, 2020). In the case of *E. gallinarum* (IGA2), it is becoming increasingly relevant as pathogens, both in hospital settings and in community-acquired infections. Usually considered of low pathogenic potential, these species are recognized as opportunistic pathogens in humans, particularly in immunocompromised individuals, and in animals, with the ability to cause severe invasive infections such as endocarditis, bacteremia, urinary tract infections, and pelvic infections (Ruoff *et al.*, 1990; Reid *et al.*, 2001; Monticelli *et al.*, 2018).

Enterobacter sp. (IGA3) showed resistance to antibiotics from the monobactam and sulfonamide classes. The *Enterobacter* genus, belonging to the Enterobacteriaceae family, is primarily associated with healthcare-associated infections. Currently, there are 22 species of *Enterobacter*, although not all are recognized to cause diseases only in humans. These species have played a significant role in the origin of various infections, such as urinary tract infections (UTIs), respiratory infections, soft tissue infections, osteomyelitis, and endocarditis, along with other clinical manifestations. These infections predominantly occur in healthcare settings, although they can also be acquired in the community (Wilberger *et al.*, 2012; Davin-Regli & Pagès, 2015).

On the other hand, *L. fusiformis* (IGA4) exhibited resistance to two antibiotics, ceftazidime and cefepime, from the cephalosporin class. This bacterial genus can be found in soil, water, and clinical environments, including hospitals. Although it is an

invasive bacterium, it is rarely associated with diseases in humans. However, isolated cases of severe sepsis due to persistent bacteremia caused by *Lysinibacillus* have been reported (Wenzler *et al.*, 2015). And, finally, *E. faecalis* (IGA5) showed no resistance to the evaluated antimicrobials. However, the relevance of this species as an opportunistic pathogen in birds and humans is highlighted, emphasizing the continued importance of monitoring and controlling antimicrobial resistance associated with *E. faecalis* (Pereira *et al.*, 2020). And some strains have been used as animal probiotics.

Research like this should be more extensively explored by other researchers, given the limited availability of data in the literature that contribute to the identification of bacteria in poultry farms. Moreover, these studies provide valuable information on management practices and appropriate treatments to ensure the health and welfare of birds.

CONCLUSIONS

This study highlighted the importance of identifying the molecular profile through the application of the 16S rRNA molecular profiling technique and evaluating antimicrobial sensitivity in bacterial isolates from poultry farms. Identifying bacteria with pathogenic potential in poultry facilities can provide crucial information for monitoring and disease control in birds. Further research is needed to perform comprehensive genotyping of bacterial isolates, including virulence and resistance gene analyses. These data play a fundamental role in prevention strategies, aiming to improve poultry health, ensure food safety, and protect both animal and public health.

ACKNOWLEDGEMENTS

We would like to thank the Biotechnology Graduate Program, the Biodiversity and Biotechnology of Bionorte Network Program, and the Vice Dean of Research of the Federal University of Tocantins (UFT). The National Council of Scientific and Technological Development (CNPq, Brazil) Project: 424435/2021-7.

Conflict of interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

BIBLIOGRAPHIC REFERENCES

ABPA. Associação Brasileira de Proteína Animal (2022). Dados e estatísticas. <https://abpa-br.org/wp-content/uploads/2023/01/abpa-relatorio-anual-2022.pdf>

Altschul, S. F.; Gish, W.; Miller, W.; Myers, E. W.; Lipman, D. J. (1990). Basic local alignment search tool. *Journal of molecular biology*. 215(3): 403-410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2)

- Arbab, S.; Ullah, H.; Wei, X.; Wang, W.; Ahmad, S. U.; Zhang, J. (2021). Drug resistance and susceptibility testing of Gram negative bacterial isolates from healthy cattle with different β -Lactam resistance Phenotypes from Shandong province China. *Brazilian Journal of Biology*. 83: e247061. <https://doi.org/10.1590/1519-6984.247061>
- Baynes, R. E.; Dedonder, K.; Kissell, L.; Mzyk, D.; Marmulak, T.; Smith, G.; Riviere, J. E. (2016). Health concerns and management of select veterinary drug residues. *Food and Chemical Toxicology*. 88: 112-122. <https://doi.org/10.1016/j.fct.2015.12.020>
- Berghiche, A.; Khenenou, T.; Labiad, I. (2018). Antibiotics resistance in broiler chicken from the farm to the table in Eastern Algeria. *Journal of World's Poultry Research*. 8(4): 95-99.
- CLSI. (2020). *Performance Standards for Antimicrobial Susceptibility Testing*. 30th ed. Wayne, PA: CLSI supplement M100. 416p.
- Cole, J. R.; Wang, Q.; Fish, J. A.; Chai, B.; McGarrell, D. M.; Sun, Y.; Tiedje, J. M. (2014). Ribosomal Database Project: data and tools for high throughput rRNA analysis. *Nucleic acids research*. 42(D1): D633-D642. <https://doi.org/10.1093/nar/gkt1244>
- Davin-Regli A.; Pagès, J. M. (2015). Enterobacter aerogenes and Enterobacter cloacae; versatile bacterial pathogens confronting antibiotic treatment. *Frontiers in microbiology*. 6: 392. [10.3389/fmicb.2015.00392](https://doi.org/10.3389/fmicb.2015.00392).
- Deusdará, T. T.; Felix, M. K. C.; Brito, H. S.; Ribeiro, D. R.; Cangussu, E. W. S.; Albuquerque, B.; Santos, G. R.; Chaves, J. R.; Carvalho, W. C. R.; Astolfi-Filho, S.; Assunção, E. N.; Mariúba, L. A. M.; Nogueira, P. A.; Viana, K. F.; Brandi, I. V.; Cangussu, A. S. R. (2023). Resistance determinants of emerging pathogens isolated from an intensive care unit as a parameter of population health conditions of the Legal Amazon microregion. *Brazilian Journal of Biology*. 83: e269778. <https://doi.org/10.1590/1519-6984.269778>
- EMBRAPA. (2021). Doenças. <https://www.embrapa.br/agencia-de-informacao-tecnologica/criacoes/frango-de-corte/producao/sanidade/doencas>
- Fardsanei, F.; Dallal, M. M. S.; Douraghi, M.; Salehi, T. Z.; Mahmoodi, M.; Memariani, H.; Nikkhahi, F. (2017). Genetic diversity and virulence genes of Salmonella enterica subspecies enterica serotype Enteritidis isolated from meats and eggs. *Microbial pathogenesis*. 107: 451-456. <https://doi.org/10.1016/j.micpath.2017.04.026>
- Gomes, A. V. S.; Silva, V. N.; Silva, M. C. (2019). Principais enfermidades em avicultura e suas implicações na qualidade do produto. *Brazilian Journal of Development*. 5(7): 12145-12157.
- Gutarowska, B.; Szulc, J.; Nowak, A.; Otłowska, A.; Okrasa, M.; Jachowicz, A.; Majchrzycka, K. (2018). Dust at various workplaces—Microbiological and toxicological threats. *International Journal of Environmental Research and Public Health*. 15(5): 877. <https://doi.org/10.3390/ijerph15050877>
- Gržinić, G.; Piotrowicz-Cieślak, A.; Klimkowicz-Pawlas, A.; Górny, R. L.; Ławniczek-Wałczyk, A.; Piechowicz, L.; Wolska, L. (2023). Intensive poultry farming: A review of the impact on the environment and human health. *Science of The Total Environment*. 858: 160014. <https://doi.org/10.1016/j.scitotenv.2022.160014>
- Hou, X. G.; Kawamura, Y.; Sultana, F.; Shu, S.; Hirose, K.; Goto, K.; Ezaki, T. (1998). Description of *Arthrobacter creatinolyticus* sp. nov., isolated from human urine. *International Journal of Systematic and Evolutionary Microbiology*. 48(2): 423-429. [10.1099/00207713-48-2-423](https://doi.org/10.1099/00207713-48-2-423)

- Kumar, S.; Stecher, G.; Li, M.; Knyaz, C.; Tamura, K. (2018). MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms. *Molecular biology and evolution*. 35(6): 1547-1549. 10.1093/molbev/msy096
- Lateef, A.; Adelere, I. A.; Gueguim-Kana, E. B (2015). The biology and potential biotechnological applications of *Bacillus safensis*. *Biologia*. 70(4): 411-419. <https://doi.org/10.1515/biolog-2015-0062>
- Magiorakos, A. P.; Srinivasan, A.; Carey, R. B.; Carmeli, Y.; Falagas, M. E.; Giske, C. G.; Monnet, D. L. (2012). Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clinical microbiology and infection*. 18(3): 268-281. 10.1111/j.1469-0691.2011.03570.x
- Manyi-Loh, C.; Mamphweli, S.; Meyer, E.; Okoh, A. (2018). Antibiotic use in agriculture and its consequential resistance in environmental sources: potential public health implications. *Molecules*. 23(4): 795. <https://doi.org/10.3390/molecules23040795>.
- Monticelli, J.; Knezevich, A.; Luzzati, R.; Di Bella, S. (2018). Clinical management of non-faecium non-faecalis vancomycin-resistant enterococci infection. Focus on *Enterococcus gallinarum* and *Enterococcus casseliflavus/flavescens*. *Journal of Infection and Chemotherapy*. 24(4): 237-246. 10.1016/j.jiac.2018.01.001
- Moyane, J. N.; Jideani, A. I. O.; Aiyegoro, O. A. (2013). Antibiotics usage in food-producing animals in South Africa and impact on human: Antibiotic resistance. *Afr. J. Microbiol. Res.* 7(24): 2990-2997. <https://doi.org/10.5897/AJMR2013.5631>
- Moretto, V. T.; Bartley, P. S.; Ferreira, V. D. M.; Santos, C. S.; Silva, L. K.; Ponce-Terashima, R. A.; Barbosa, L. M. (2021). Microbial source tracking and antimicrobial resistance in one river system of a rural community in Bahia, Brazil. *Brazilian Journal of Biology*. 82: e231838. <https://doi.org/10.1590/1519-6984.231838>
- Paudel, D.; Boogaard, H.; de Wit, A.; Janssen, S.; Osinga, S.; Pylaniadis, C.; Athanasiadis, I. N. (2021). Machine learning for large-scale crop yield forecasting. *Agricultural Systems*. 187: 103016. <https://doi.org/10.1016/j.agsy.2020.103016>.
- Pereira, V. C.; Romero, L. C.; Pinheiro-Hubinger, L.; Oliveira, A.; Martins, K. B.; Cunha, M. L. R. S. (2020). Coagulase-negative staphylococci: a 20-year study on the antimicrobial resistance profile of blood culture isolates from a teaching hospital. *Brazilian Journal of Infectious Diseases*. 24: 160-169. 10.1016/j.bjid.2020.01.003
- Reid K. C.; Cockerill III F. R.; Patel R. (2001). Clinical and Epidemiological Features of *Enterococcus casseliflavus/flavescens* and *Enterococcus gallinarum* Bacteremia: A Report of 20 Cases. *Clinical Infectious Diseases*. 32: 1540- 1546. 10.1086/320542
- Ruoff, K. L.; De La Maza, L.; Murtagh, M. J.; Spargo, J. D.; Ferraro, M. J. (1990). Species identities of enterococci isolated from clinical specimens. *Journal of Clinical Microbiology*. 28(3): 435-437. 10.1128/jcm.28.3.435-437.1990
- Sanger, F.; Nicklen, S.; Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors. *Proceedings of the National Academy of Sciences*. 74(12): 5463-5467. 10.1073/pnas.74.12.5463
- Santos, R. G.; Hurtado, R.; Gomes, L. G. R.; Profeta, R.; Rifici, C.; Attili, A. R.; Azevedo, V. (2020). Complete genome analysis of *Glutamicibacter creatinolyticus* from mare abscess and comparative genomics provide insight of diversity and adaptation for *Glutamicibacter*. *Gene*. 741: 144566. 10.1016/j.gene.2020.144566

- Wenzler, E.; Kamboj, K.; Balada-Llasat, J. M (2015). Severe sepsis secondary to persistent *Lysinibacillus sphaericus*, *Lysinibacillus fusiformis* and *Paenibacillus amylolyticus* bacteremia. *International Journal of Infectious Diseases*. 35: 93-95. 10.1016/j.ijid.2015.04.016
- Wilberger, M. S.; Anthony, K. E.; Rose, S. McClain, M.; Bermudez, L. E. (2012). Beta-lactam antibiotic resistance among *Enterobacter* spp. isolated from infection in animals. *Advances in Microbiology*. 2: 129-137. 10.4236/aim.2012.22018
- Zarrilli, R.; Pournaras, S.; Giannouli, M.; Tsakris, A. (2013). Global evolution of multidrug-resistant *Acinetobacter baumannii* clonal lineages. *Int. J. Antimicrob. Agents*. 41(1): 11-19. <https://doi.org/10.1016/j.ijantimicag.2012.09.008>