

e-ISSN Online 2256-2273

Received: October 29 2024

Accepted: March 11 2025

Published: April 28 2025

*Research article: Agriculture* 

# Resistance of agricultural pests to synthetic insecticides in Colombia: A review

Resistencia de plagas agrícolas a insecticidas sintéticos en Colombia: Artículo de revisión

Maria Pineda<sup>1</sup>; Tito Bacca<sup>2</sup>

University of Reggio Calabria, Reggio Calabria, Italia, maria.pineda@unirc.it, https://orcid.org/0000-0002-7947-3278
 Universidad del Tolima, Ibagué, Tolima, Colombia, titobacca@ut.edu.co, https://orcid.org/0000-0002-2960-5527

Cite: Pineda, M.; Bacca, T. (2025). Resistance of agricultural pests to synthetic insecticides in Colombia: A review. *Revista de Ciencias Agrícolas*. 42(1): e1252. https://doi.org/10.22267/rcia.20254201.252

## ABSTRACT

The phenomenon of insecticide resistance is defined as the ability of insect populations to develop the capacity to survive doses of insecticides that would otherwise be lethal. This phenomenon can be attributed to the natural selection of individuals with genetic mutations that confer resistance. A comprehensive understanding of insecticide resistance mechanisms is crucial for developing novel control strategies within existing integrated pest management plans. In this context, this review examines the research conducted to characterize the resistance of agricultural pests (insects/mites) to insecticides/acaricides in Colombia. A search of relevant scientific literature was conducted using keywords in the databases. A literature review revealed that, according to reports from Colombia, there have been 27 documented instances of insecticide/acaricide resistance across 12 species. Conversely, the Arthropod Pesticide Resistance Database (APRD) indicates the existence of 98 reports of resistance to insecticides/acaricides distributed across 10 species. A single instance of acaricide resistance was identified in the respiratory target; all other cases demonstrated resistance to the nervous and muscular targets. In this review, we present the main results, which are focused on determining toxicity curves and calculating lethal concentrations. We highlight the research carried out in the last decade, which makes use of new molecular techniques and allows a more detailed view of the resistance mechanisms. Furthermore, we suggest novel

**Keywords:** chemical control; crops pest; insecticide; integrated pest management; resistance; toxicology

# RESUMEN

La resistencia a insecticidas es un fenómeno en el cual las poblaciones de insectos desarrollan la capacidad de sobrevivir a dosis de insecticidas que normalmente serían letales. Esto debido a la selección natural de individuos con mutaciones genéticas que les confieren la resistencia. El conocimiento de los mecanismos de resistencia a los insecticidas es indispensable para reformular las estrategias de control en los actuales planes de manejo integrado de plagas. En este contexto, esta revisión analizó las investigaciones que tuvieron como objetivo caracterizar la resistencia de plagas agrícolas (insectos/ácaros) a insecticidas/acaricidas en Colombia. Utilizamos palabras claves para realizar la búsqueda de todas las investigaciones científicas relacionadas en bases de datos. El análisis de la información demostró que, en Colombia, existen 27 reportes de resistencia a insecticidas/ acaricidas distribuidos en 12 especies. Por otro lado, la base de datos sobre resistencia a plaguicidas de artrópodos (APRD) relata la existencia de 98 reportes distribuidos en 10 especies. Encontramos que únicamente existe un reporte de un acaricida resistente al modo de acción respiratorio. Los demás reportes presentan resistencia al modo de acción del sistema nervioso y muscular. En esta



revisión presentamos los principales resultados que están enfocados en la determinación de curvas de toxicidad y cálculo de concentraciones letales. Destacamos las investigaciones realizadas en la última década, las cuales hacen uso de nuevas técnicas moleculares y permiten una visión detallada de los mecanismos de resistencia. Finalmente, planteamos algunas sugerencias para incluir metodologías novedosas que permitan mejorar los actuales planes de manejo integrado de plagas.

**Palabras clave:** control químico; insecticida, manejo integrado de plagas; plagas de cultivos; resistencia; toxicología

#### **INTRODUCTION**

For several decades, the use of agrochemicals for control of pests has been the most prevalent method globally. As a result, farmers have resorted to the indiscriminate use of broad-spectrum synthetic pesticides, often unregistered and applied in excessive doses. Consequently, pests have developed resistance to most major pesticide classes. Moreover, the widespread use of these toxic, nonselective pesticides has led to harmful impacts on the health of users, consumers, and non-target organisms, including pollinators and natural predators of insect pests (Cutler & Guedes, 2017; Guedes et al., 2016). Insect resistance to insecticides is defined by IRAC as "an inherited change in the susceptibility of a pest population which is reflected in the repeated failure of an insecticide to achieve the expected level of control when used according to label recommendations for that pest species." This phenomenon is very common due to the misuse of insecticides, mainly because of the selection pressure exerted by the same mode of action of an insecticide on a target pest (IRAC, 2024a). The first documented case of resistance was in 1914; since then, the numbers have been increasing to 16,570 arthropods resistant to at least one class of pesticide have been reported. including mites and insects (Sparks et al., 2020).

In Colombia, the agricultural sector plays a crucial role in the country's economy and food security. The issue of insecticide resistance represents a significant challenge, particularly given the limited research conducted in Colombia. This is due to a lack of interest and awareness regarding its importance in integrated pest management plans. However, there is a greater body of knowledge available on insect vectors of tropical diseases that are medically important. This is because, in 2004, the National Institute of Health of Colombia established the Network for Surveillance of Resistance to Insecticides Used in Public Health. Consequently, there is more extensive documentation of mosquito (Culicidae) resistance to the use of insecticides (Aguirre-Obando *et al.*, 2015; Love *et al.*, 2023; Orjuela *et al.*, 2018).

According to the Arthropod Pesticide Resistance Database (APRD) (Mota-Sanchez & Wise, 2024), there have been 199 documented instances of resistance in Colombia. Of these, 51% pertain to arthropods of medical and veterinary importance. This category includes eight species (101 reports) of mites (Acari) and mosquito vectors of diseases (Diptera: Culicidae). In contrast, the remaining 49% pertains to agricultural importance, distributed among 10 species with 98 reports. This encompasses one species of phytophagous mite and nine species of insects, with Hemiptera and Lepidoptera being particularly prevalent. Some reports suggest that most information on resistance in Colombia relates to *Helicoverpa virescens* in cotton crops and the whitefly complex in solanaceous and flower crops (Rodríguez & Cardona, 2001).

Currently, there are no research programs in Colombia, and the number of resistant insects and resistance mechanisms remain unknown. Thus, our objective



was to conduct the first literature review to obtain a clearer picture of the current situation in our country. It is important to note that insecticide resistance is, and will continue to be, one of the reasons driving the discovery of new molecules for chemical pest control (Lamberth *et al.*, 2013; Maienfisch & Stevenson, 2015; Sparks & Lorsbach, 2017). Nevertheless, the deployment of insecticides represents one of the fundamental pillars of Integrated Pest Management (IPM). It is therefore recommended to establish research lines in the field of agricultural toxicology to evaluate control failures and develop new criteria to modify current management plans while incorporating more sustainable and environmentally friendly alternatives in fumigant applications.

#### Search strategy

An exhaustive search of bibliographic material in national and international databases was conducted to extract data on the resistance of agricultural pests to insecticides/acaricides in Colombia. No restrictions were placed on the date or type of publication of the information. Articles about resistance and toxicology in the context of medical and veterinary entomology were excluded from the review. A total of 27 documents were obtained, distributed as follows: 16 articles published in national and international journals, nine abstracts and proceedings from national congresses related to agricultural entomology, and two degree theses. Consequently, a total of 17 arthropod species in Colombia have demonstrated resistance to at least one pesticide (Table 1).

Taxonomy	Species	No. Reports	References	
Acari: Tetranychidae	<i>Tetranychus cinnabarinus</i> (Boisduval, 1867)	2a (Murillo & Mosquera, 1984)		
	<i>Tetranychus bimaculatus</i> Harvey, 1892	1b	(Mota-Sanchez & Wise, 2024)	
Lepidoptera: Noctuidae	<i>Heliothis virescens</i> Fabricius, 1777	<i>eliothis virescens</i> abricius, 1777 5a (Lozano Cruz, 196 1977; Rendón <i>et a</i> <i>et al.</i> , 1993; Wolfe 1973)		
	<i>Heliothis spp.</i> Ochsenheimer, 1816	4a	(Alcaraz, 1971; Collins, 1987; Rendon <i>et al.</i> , 1978; Rendon & Cardona, 1976)	
	<i>Heliothis zea</i> Boddie, 1850	2b	(Mota-Sanchez & Wise, 2024)	
	<i>Helicoverpa zea</i> Boddie, 1850	1b	(Mota-Sanchez & Wise, 2024)	
	<i>Spodoptera frugiperda</i> J. E. Smith, 1797	2a - 2b	(Mota-Sanchez & Wise, 2024; Ríos- Díez & Saldamando-Benjumea, 2011; Zenner de Polanía, 1996)	
Lepidoptera: Gelechiidae	Tecia solanivora (Povolný, 1973)	2a - 4b	(Bacca <i>et al.</i> , 2017; Gutiérrez <i>et al.,</i> 2019; Mota-Sanchez & Wise, 2024)	
	Tuta absoluta (Meyrick, 1917)	1b	(Haddi <i>et al.</i> , 2012)	
Lepidoptera: Crambidae	<i>Ostrinia nubilalis*</i> Hübner,1796	1b	(Mota-Sanchez & Wise, 2024)	

#### **Table 1.** List of species of arthropods resistant to pesticides in Colombia



Lepidoptera: Erebidae	Alabama argillacea (Hübner, 1823)	1b (Mota-Sanchez & Wise, 2024)		
Hemiptera: Aleyrodidae	Bemisia tabaci (Gennadius, 1889)	3a - 23b	(Mota-Sanchez & Wise, 2024; Rodríguez <i>et al.,</i> 2005, Rodríguez <i>et al.,</i> 2012)	
	Trialeurodes vaporariorum (Westwood, 1856)	2a - 59b	(Buitrago <i>et al.,</i> 1994; Mota-Sanchez & Wise, 2024; Rodríguez & Cardona, 2001)	
Coleoptera: Curculionidae (Scolitynae)	Hypothenemus hampei (Ferrari, 1867)	1a	(Navarro <i>et al.,</i> 2010)	
Coleoptera: Bostrichidae	<i>Rhyzopertha dominica</i> (Fabricius, 1792)	1a	(Ortega <i>et al.,</i> 2021)	
Coleoptera: Chrysomelidae (Bruchinae)	Zabrotes subfasciatus (Bohemann, 1833)	1a - 1b	(Mota-Sanchez & Wise, 2024; Tyler & Evans, 1981)	
	<i>Thrips palmi</i> Karny, 1925	18	(Rodríguez <i>et al.,</i> 2003)	

Pineda & Bacca - Resistance agricultural pests to synthetic insecticides

**a.** Number of reports in the literature. **b.** Number of reports in Database APRD (Mota-Sanchez & Wise, 2024). https://www.pesticideresistance.org/.\*. According to the European and Mediterranean Plant Protection Organization (EPPO) Global Database (2024) and a list of quarantine pests that are absent in Colombia, as provided by the Colombian Agricultural Institute, ICA (2024), O. *nubilalis* has not been reported in Colombia.

Among insects, the order Lepidoptera is of particular note (60%), with the greatest quantity of resistance information reported in the genus *Helicoverpa* spp. (formerly known as *Heliothis*), which is an important defoliator of cotton and corn crops. In the Coleoptera order (20%), certain pests of stored grains and seeds merit particular attention, while in the Hemiptera order (13%), whiteflies — insects associated with virus transmission — warrant further investigation. Finally, the order Thysanoptera reports one species (7%). Additionally, two species of the mite genus *Tetranychus*, which are known to be phytophagous pests, have been identified as resistant to acaricides in multiple crops (Figure 1).





The veracity of the above-listed pests was corroborated through a review of databases maintained by the Colombian Agricultural Institute (ICA, 2024), the National Open Data Network on Biodiversity (SIB, 2024), and the European and Mediterranean Plant Protection Organization, EPPO Global Database (EPPO, 2024). The objective of this review was to ascertain the presence of these pests within the Colombian context. Our comprehensive analysis revealed that *O. nubilalis*, a European lepidopteran commonly known as the corn borer, is included in the list of absent quarantine pests that is updated annually by the Colombian Agricultural Institute (ICA, 2024). However, despite its inclusion in the Arthropod Pesticide Resistance Database (APRD) of Mota-Sanchez & Wise (2024), there is a dearth of studies investigating insecticide resistance in this pest within the Colombian context. We attribute this absence to an error in the APRD database.

The use of insecticides is a common practice for controlling insect pests in a variety of crops. After the Second World War and the successful introduction of dichlorodiphenyltrichloroethane (DDT), mainly for controlling diseasetransmitting insects, it was banned due to its persistent nature and the many contamination problems it caused (Beard, 2006; Jarman & Ballschmiter, 2012; Sánchez-Bayo, 2019). Later, alternative synthetic insecticides emerged, including organophosphates and carbamates (Gupta *et al.*, 2022), which were used as broad-spectrum insecticides due to their ease of acquisition. In Colombia, they were used excessively and with inadequate practices in potato, cotton, corn, and flower crops. Due to the lack of other insecticides with a different mode of action, pests such as the whitefly complex quickly became resistant.

Undoubtedly, the advent and introduction of pyrethroids in the 1980s provided a viable and successful alternative for controlling many pests. In Colombia, they were successfully introduced to control H. virescens and other Lepidoptera in cotton crops. However, due to the lack of alternatives, repeated use and improper dosage in Tolima (Colombia) resulted in control failures and the selection of populations resistant to pyrethroids and organophosphates (Rendon et al., 1977; Rendon et al., 1978; Rendon & Cardona, 1976). Additionally, the corn earworm, S. frugiperda, was reported to exhibit resistance to the same groups in 1985, including carbamates (Table 2) (Zenner de Polanía, 1996). In the 1990s, it was the whitefly complex, consisting of *B. tabaci* and *T. vaporariorum*, which are important pests in solanaceous and flower crops. Additionally, the whitefly was found to be associated with virus transmission and demonstrated resistance to the same groups, as well as to organochlorines and neonicotinoids. This finding was based on a synthesis of multiple studies conducted over several years by the International Center for Tropical Agriculture (CIAT) (Buitrago et al., 1994; Rodríguez et al., 2005).

In subsequent years, several studies have reported resistance to the same mode of action across different chemical groups: pyrethroids in *T. absoluta*, spinosyns, carbamates and organophosphates in *T. palmi*, and organochlorines in *Z. subfasciatus* and *H. hampei* (Table 2). In the last decade, resistance to permethrin, chlorpyrifos, and carbofuran has been reported in *T. solanivora* and to pyrethroids in *R. dominica*. Thus, 26 studies report resistance to insecticides with modes of action targeting nerve and muscle systems. Finally, in Colombia, there is only one record of resistance to the respiration-target mode of action of Cihexatina in the mite *T. cinnabarinus*. A mitochondrial respiration produces ATP, the molecule that energizes all vital cellular processes. The insecticide inhibits the enzyme that synthesizes ATP (IRAC, 2024b).



It is estimated that 98% of the reported cases show resistance at the target site, but these studies are not sufficient to better understand the molecular and genetic basis of resistance mechanisms (Bass & Nauen, 2023; Sajad *et al.*, 2020)

Mode of action MoA *	No.	Sub-group	Active Ingredient	Pest
<b>NERVE AND</b> <b>MUSCLE TARGETS</b> Acetylcholinesterase (AChE) inhibitors Inhibit AChE, causing hyperexcitation. AChE is the enzyme that terminates the action of the excitatory neurotransmitter acetylcholine at nerve synapses.		Carbamates	Carbofuran	Tecia solanivora
			Carbosulfan	Thrips palmi
	1A		Methomyl	Bemisia tabaci Spodoptera frugiperda Trialeurodes vaporariorum
		Organophosphates	Chlorpyrifos	Spodoptera frugiperda; Tecia solanivora
			Methamidophos	Bemisia tabaci Trialeurodes vaporariorum
			Monocrotophos	Trialeurodes vaporariorum
	1B		Omethoate	Tetranychus cinnabarinus
			Parathion Methyl	Heliothis virescens
			Pirimiphos methyl	Rhyzopertha dominica
			Profenofos	Trialeurodes vaporariorum
			Triazophos	
			Toxametil	Heliothis virescens
			(Methamidophos)	
NERVE AND MUSCLE TARGETS GABA-gated chloride channel blockers Block the GABA-activated chloride channel, causing hyperexcitation and convulsions. GABA is the major inhibitory neurotransmitter in insects.	2A	Cyclodiene: organochlorines	Endosulfan	Heliothis virescens
		Organochlorine	Dieldrin	Hypotenemus hampei
			Endrin	Heliothis virescens
	NC		Thiocyclam hydrogen oxalate	Bemisia tabaci
			Dienochlor	Tetranychus cinnabarinus
			Gamma-HCH (Lindano)	Zabrotes subfasciatus

Table 2. Classification of insecticides that are registered for resistance in Colombia



Mode of action MoA *	No.	Sub-group	Active Ingredient	Pest
<b>NERVE AND</b> <b>MUSCLE TARGETS</b> Sodium channel modulators Keep sodium channels open, causing hyperexcitation and, in some cases, nerve block. Sodium channels are		Pyrethroids	Bifenthrin	Bemisia tabaci Rhyzopertha dominica
			Cypermethrin	Bemisia tabaci Heliothis virescens Spodoptera frugiperda Trialeurodes vaporariorum
	3A		Deltamethrin	Heliothis virescens Trialeurodes vaporariorum Rhyzopertha dominica
involved in the propagation of action potentials along			Fenvalerate	Heliothis virescens
nerve axons.			Lambda-Cyhalothrin	Spodoptera frugiperda
			Permethrin	Tecia solanivora
	3B	DDT	DDT	Heliothis virescens
NERVE AND MUSCLE TARGETS Nicotinic acetylcholine receptor (nAChR) competitive modulators Bind to the acetylcholine site on nAChRs, causing a range of symptoms from hyper-excitation to lethargy and paralysis. Acetylcholine is the major excitatory neurotransmitter in the insect central nervous system.	4A	Neonicotinoids	Imidacloprid	Bemisia tabaci Thrips palmi
			Thiamethoxam	Bemisia tabaci
NERVE AND MUSCLE TARGETS Nicotinic acetylcholine receptor (nAChR) allosteric modulators - Site I: Allosterically activate nAChRs, causing hyperexcitation of the nervous system. Acetylcholine is the major excitatory neurotransmitter in the insect central nervous system.	5	Spinosyns	Spinosyns	Thrips palmi





\*IRAC: Mode of action classification scheme, version 11.2, 08/2024; NC: No current classification

#### Mechanisms of resistance

As previously stated, resistance is defined as the ability of an insect population to survive and successfully reproduce following exposure to a dose of insecticide that was previously effective in controlling it (Sparks & Nauen, 2015). However, this phenomenon of insecticide resistance is a natural process and is part of the natural process of evolution and adaptation (Costantini, 2019; Madgwick *et al.*, 2023). This adaptive phenomenon involves a strong selection of specific mutations that confer different types of resistance, thus causing failures in IPM (IRAC, 2024a).

However, this phenomenon of insecticide resistance is a natural occurrence and an integral aspect of the evolutionary process (Madgwick et al., 2023). This adaptive phenomenon involves the strong selection of specific mutations that confer different types of resistance, leading to failures in IPM. It is unfortunate that in many cases, resistance is not only present to a chemical compound but in some cases, it also confers cross-resistance to other structurally related agents. This is due to the structural similarity of the compounds and the fact that in many cases, they can share a common target site within the pest, such that they share a common mechanism of action (IRAC, 2024a; Sajad et al., 2020). The hereditary changes present in a pest are associated with genetic mutations that in turn can take different forms and cause different types of resistance, according to IRAC (2024a): a. Target Site Resistance: The occurrence of a mutation at a specific site on the receptor protein results in a reduction in the efficacy of the insecticide (Constant, 1999). **b. Metabolic resistance:** In this case, insects develop the ability to break down the insecticide through the action of enzymes. For example, they produce higher levels of detoxifying enzymes (such as esterases, cytochrome P450 monooxygenases, or glutathione S-transferases) that can break down or alter the insecticide before it reaches its target site. Additionally, the insect's metabolism adapts to degrade the insecticide more rapidly, reducing its effective concentration (Ranganathan et al., 2022). c. Physical adaptation: In this case, the insect develops a physical adaptation that enables it to evade the insecticide, such as a thicker cuticle, an additional wax layer, or accelerated waste excretion. These adaptations, on their own, do not provide significant protection. However, they are often combined with other mechanisms to enhance effectiveness (Perry et al., 2011). d. Behavioral adaptation: This particular form of resistance is less prevalent, as it entails mutations that have altered the insect's intrinsic behavioral patterns. In order to evade contact with the insecticide, insects may modify their behavior, which may include alterations to their feeding, mating, or shelter-seeking habits, thereby reducing their exposure (Zalucki & Furlong, 2017).

Insects can develop resistance to multiple insecticides, often due to a combination of the above mechanisms or because one mechanism (e.g., detoxification) may



confer resistance to insecticides of different chemical classes. This phenomenon is referred to as multiple and cross-resistance. Consequently, the classification of modes of action established by IRAC is intended to assist growers in identifying distinct modes of action, thereby preventing the repetitive utilization of analogous insecticide products and averting the development of resistance (Gnanadhas *et al.*, 2013; Sparks & Nauen, 2015).

# Study of resistance in Colombia

The most prevalent form of insecticide resistance is the result of genetic mutations at the site of action, which impair the efficacy of the insecticide. The common examples of this phenomenon include insecticides with a target mode of action in the nervous and muscular systems. A clear example is provided by mutations in the enzyme acetylcholinesterase, which confers resistance to organophosphates, or in voltage-dependent sodium channels, which confers resistance to pyrethroids (Chrn *et al.*, 2016; Liu, 2012). In our extensive review, we found that 98% of the literature reports resistance to this mode of action, with multiple species exhibiting resistance to pyrethroids, organophosphates, and carbamates. The remaining two percent can be attributed to the respiration targets' mode of action (Table 2; Figure 2). Of the published papers (n=16), 75% of the research is based on basic toxicity tests using the impregnated vial technique (IRAC, 2011), direct application on immature stages, immersion of the arthropods in different concentrations of the pesticides studied, or in some more specific cases, immersion of the foliage for sucking pests.





Each methodology was adapted to the specific conditions of each laboratory and the experience of the researchers involved. All of the methodologies included pre-tests to determine a mortality rate and subsequent calibration curves to determine the new lethal concentrations (LC) of each product used. Concentration values are calculated using Probit analysis and Abbott's mortality correction. In the majority of studies, comparisons are made between the LC values of a susceptible and a resistant population to determine the resistance ratio (RR) of the species under research to a given compound (Akçay, 2013; Sakuma, 1998). These studies allow the deduction of whether a population is resistant to a specific compound due to the influence of selection pressure. Nevertheless, these studies are unable to ascertain the physiological resistance mechanism that the pest has acquired. It is crucial to study the biological, genetic, and molecular bases of insecticide resistance because it allows for the determination of which physiological process inactivates the insecticide before it reaches its molecular target (Bass & Nauen, 2023; Perry *et al.*, 2015).

A review of the Colombian literature revealed that only four publications (25%) employed methodologies that facilitate a more comprehensive understanding of resistance mechanisms. One study by Valencia et al. (1993) involved enzymatic activity analyses, including esterases, carboxylesterases, the mixed-function oxidase system (MFOs), and cytochrome P-450 activity. This analysis demonstrated an increase in the enzymatic activity of the resistant population of H. virescens to Triazophos. Newer techniques such as highresolution fusion polymerase chain reaction (PCR) quantification to identify polymorphisms (SNPs) and mutations for rapid identification of Rdl (resistance to dieldrin) alleles in different pest populations were used by Navarro et al. (2010), where they reported the Rdl allele in populations of *H. hampei*. Similar studies identified pyrethroid resistance mediated by mutation of the para-type sodium channel in various populations of *T. absoluta*. Haddi et al. (2012) showed that the Rio Negro (Antioquia-Colombia) population exhibited mutations in dr/ super-kdr type genes (M918T, T929I and L1014F), which attribute resistance to pyrethroids. Similar results were reported by Bacca et al. (2017), who determined genetic mutations of Kdr genes (L1014F) conferring pyrethroid resistance in T. solanivora of Nariño and Boyacá departments populations.

#### Challenges and perspectives of resistance in Colombia

Studies on the resistance of agricultural pests to insecticides in Colombia are insufficient due to a lack of knowledge of their importance in IPM programs. In addition to research on the biology of resistant insects that allows us to determine the adaptive costs of the species, it is important to include methodologies related to genetic, biochemical, and molecular analyses to understand the metabolic bases, resistance mediated by the target site, and identify new insecticide targets, in addition to understanding the adaptive costs of resistant species and mitigating the impact currently (Erdogan *et al.*, 2024; Rix *et al.*, 2022; Wang & Wang, 2024). In contrast to surveillance programs for insect vectors of diseases, in Colombia, there are no research and monitoring programs in insecticide toxicology to detect failures in control plans in time and create new strategies that are effective and specific for each species.

General recommendations to avoid and reduce control failures focus on rotation plans for different types of insecticide/acaricide modes of action, using the doses recommended by the manufacturer, respecting the frequency and, if possible, using low-persistence insecticides, taking into account the collateral effect on beneficial insects. In Colombia, some political and legal measures are promoted for specific quarantine pests, regulated by the Colombian Agricultural Institute (ICA). Despite the introduction of new integrated management strategies for specific crops and pests to reduce the impact of resistance, the challenge is difficult. There is a great deal of information on resistance to various insecticides in the carbamate, organophosphate, and pyrethroid groups (Table 2) for *H. virescens* in cotton and *S. frugiperda* in maize and rice, but integrated management practices that have rationalized insecticide use have in some cases restored susceptibility, for example, in corn, with the introduction of new molecules such as diflubenzuron and Bt crops. In 2002, transgenic maize and cotton crops were introduced, containing proteins (Cry1Ab or Cry1F) of Bacillus thuringiensis (Bt), designed to reduce resistant populations and the use of



synthetic insecticides in pest control (Blanco *et al.*, 2016; Monnerat *et al.*, 2006). In 2018, the University of Tolima registered the susceptibility of *S. frugiperda* to lambda-cyhalothrin and methomyl, attributed to the use of BT maize crops (partial findings have been disseminated SOCOLEN congress) (Jaramillo-Barrios *et al.*, 2020; Ríos-Díez & Saldamando-Benjumea, 2011; Rodriguez-Chalarca *et al.*, 2024; Valencia-Cataño *et al.*, 2016; Zenner *et al.*, 2005).

However, there are also many examples of resistance that are challenging, such as *T. solanivora* in potato crops, where resistant populations have changed their biology as part of their adaptive process. Another example is the whitefly complex, which is highly accentuated in southern Colombia owing to the indiscriminate use of insecticides banned in Colombia but easily accessible due to access to neighboring countries where they are marketed. In this case, the inclusion of new public policies in the agricultural sector is necessary to improve the extension service and attention to farmers, which will make it possible to diagnose pests classified as resistant and create a monitoring network for potentially resistant pests.

The agricultural extension and assistance service in Colombia is primarily managed by personnel from multinational corporations that produce, market, and distribute pesticides. Consequently, there is a dearth of deliberate technical assistance that is genuinely oriented towards integrated pest management plans. Instead, the emphasis is on increasing sales for commissions. A considerable number of products are recommended despite lacking ICA registration for the crop and for the pests and are also used in excess. For this reason, there are control failures in many crops such as avocado, flowers, ornamentals, tomatoes, potatoes, cape gooseberry, and grapes.

Resistance and toxicology studies should go beyond the calculation of new LC50 and toxicity curves; they should aim to determine the metabolic mechanisms used by the pest to detoxify the lethal effect of the insecticide. New molecular tools should be used to analyze in detail the mutations of the genome of each species and to carry out complementary studies on sublethal effects, and adaptive and behavioral costs, to implement specific integrated pest management plans that also include environmentally friendly and sustainable practices.

## CONCLUSIONS

The evolution of pest resistance to synthetic insecticides is an evolutionary process driven by selection pressure, which is exacerbated by their inappropriate use. The high reproductive rate of insects, their enzymatic detoxification mechanisms, and selective pressure are all factors that contribute to the proliferation of individuals that possess resistant genes. Addressing this issue necessitates further research to accurately identify resistance mechanisms and formulate species-specific IPM strategies.



## REFERENCES

- Aguirre-Obando, O. A.; Bona, A. C. D.; Duque L. J. E.; Navarro-Silva, M. A. (2015). Insecticide resistance and genetic variability in natural populations of *Aedes* (Stegomyia) *aegypti* (Diptera: *Culicidae*) from Colombia. *Zoologia* (Curitiba). 32(1): 14–22. https://doi.org/10.1590/S1984-46702015000100003
- Akçay, A. (2013). The Calculation of LD50 Using Probit Analysis. *The FASEB Journal*. 27(1): 1217-28. https://doi.org/10.1096/fasebj.27.1\_supplement.1217.28
- Alcaraz, V. H. (1971). *Probables problemas de control de plagas que puedan presentarse en el cultivo del algodón en Colombia y sus posibles soluciones.* 1. Edición. Bogotá, D.C.: Federación Nacional de Algodoneros. 20p.
- Bacca, T.; Haddi, K.; Pineda, M.; Guedes, R. N. C.; Oliveira, E. E. (2017). Pyrethroid resistance is associated with a kdr -type mutation (L1014F) in the potato tuber moth *Tecia solanivora*. *Pest Management Science*. 73(2): 397–403. https://doi.org/10.1002/ps.4414
- Bass, C.; Nauen, R. (2023). The molecular mechanisms of insecticide resistance in aphid crop pests. *Insect Biochemistry and Molecular Biology*. 156: 103937. https://doi.org/10.1016/j. ibmb.2023.103937
- Beard, J. (2006). DDT and human health. *Science of the total environment*. 355(1–3): 78–89. https://doi.org/10.1016/j.scitotenv.2005.02.022
- Blanco, C.; Chiaravalle, W.; Dalla-Rizza, M.; Farias, J.; García-Degano, M.; Gastaminza, G.; Mota-Sánchez, D.; Murúa, M.; Omoto, C.; Pieralisi, B.; Rodríguez, J.; Rodríguez-Maciel, J.; Terán-Santofimio, H.; Terán-Vargas, A.; Valencia, S.; Willink, E. (2016). Current situation of pests targeted by Bt crops in Latin America. *Current Opinion in Insect Science*. 15: 131–138. https:// doi.org/10.1016/j.cois.2016.04.012
- Buitrago, N. A.; Acosta, A.; Cardona, C. (1994). Niveles de resistencia a insecticidas en *Trialeurodes vaporariorum* (Westwood) (Homoptera: Aleyrodidae), plaga del fríjol común. *Revista Colombiana de Entomología*. 20(2): 108–114.
- Chrn, C.; Shi, X.; Gao, X. (2016). Mechanism of insect metabolic resistance to pyrethroid insecticides. *Chinese Journal of Pesticide Science*. 18(5): 545–555. https://doi.org/10.16801/j.issn.1008-7303.2016.0078
- Collins, M. D. (1987). Manejo de la resistencia en *Heliothis spp* a los insecticidas piretroides. https://doi.org/http://hdl.handle.net/20.500.12324/13701
- Constant, R. (1999). Target site mediated insecticide resistance: what questions remain? *Insect Biochemistry and Molecular Biology*. 29(5): 397–403. https://doi.org/10.1016/S0965-1748(99)00024-7
- Costantini, D. (2019). Hormesis Promotes Evolutionary Change. *Dose-Response*. 17(2): 155932581984337. https://doi.org/10.1177/1559325819843376
- Cutler, G. C.; Guedes, R. N. C. (2017). Occurrence and Significance of insecticide-induced hormesis in insects. In: Duke, S. O.; Kudsk, P.; Solomon, K.(eds). *Pesticide Dose: Effects on the Environment and Target and Non-Target Organisms*. pp. 101–119. 1st edition. Washington, DC: Agricultural Research Service. 197p. https://doi.org/10.1021/bk-2017-1249.ch008
- EPPO. (2024). European and Mediterranean Plant Protection Organization Organisation Européenne et Méditerranéenne pour la Protection des Plantes. https://www.eppo.int/
- Erdogan, C.; Toprak, U.; Gurkan, M. O. (2024). Biochemical and molecular analyses of insecticide resistance in greenhouse populations of *Bemisia tabaci* (Hemiptera: Aleyrodidae) in Türkiye. *Phytoparasitica*. 52(2): 41. https://doi.org/10.1007/s12600-024-01155-5
- Gnanadhas, D. P.; Marathe, S. A.; Chakravortty, D. (2013). Biocides resistance, cross-resistance mechanisms and assessment. *Expert Opinion on Investigational Drugs*. 22(2): 191–206. https://doi.org/10.1517/13543784.2013.748035
- Guedes, R. N. C.; Smagghe, G.; Stark, J. D.; Desneux, N. (2016). Pesticide-Induced Stress in Arthropod Pests for Optimized Integrated Pest Management Programs. *Annual Review of Entomology*. 61(1): 43–62. https://doi.org/10.1146/annurev-ento-010715-023646
- Gupta, R. C.; Doss, R. B.; Yurdakok-Dikmen, B.; Malik, J. K.; Zaja-Milatovic, S.; Milatovic, D. (2022). Organophosphates and carbamates. In: Gupta, R. *Reproductive and Developmental*



*Toxicology*. pp. 617–639. 3th Edition. Academic Press. 1460p. https://doi.org/10.1016/B978-0-323-89773-0.00033-3

- Gutiérrez, Y.; Bacca, T.; Zambrano, L. S.; Pineda, M.; Guedes, R. N. (2019). Trade-off and adaptive cost in a multiple-resistant strain of the invasive potato tuber moth *Tecia solanivora*. Pest Management Science. 75(6): 1655–1662. https://doi.org/10.1002/ps.5283
- Haddi, K.; Berger, M.; Bielza, P.; Cifuentes, D.; Field, L. M.; Gorman, K.; Rapisarda, C.; Williamson, M. S.; Bass, C. (2012). Identification of mutations associated with pyrethroid resistance in the voltage-gated sodium channel of the tomato leaf miner (*Tuta absoluta*). *Insect Biochemistry and Molecular Biology*. 42(7): 506–513. https://doi.org/10.1016/j.ibmb.2012.03.008
- ICA. (2024). Reglamentación de Plagas. Plagas Cuarentenarias Ausentes. https://www.ica.gov.co/ areas/agricola/servicios/epidemiologia-agricola/plagas-reglamentadas/plagas-cuarentenariasausentes
- IRAC. (2011). Susceptibility Test Methods Series. 1, 1-2. https://irac-online.org/test-methods/
- IRAC. (2024a). Mechanisms resistance. https://irac-online.org/training-centre/resistance/ mechanisms/
- IRAC. (2024b). Test Method Library. https://irac-online.org/test-methods/test-method-library/
- Jaramillo-Barrios, C. I.; Varón-Devia, E. H.; Monje-Andrade, B. (2020). Economic injury level and action thresholds for *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) in maize crops. *Revista Facultad Nacional de Agronomía Medellín.* 73(1): 9065–9076. https://doi. org/10.15446/rfnam.v73n1.78824
- Jarman, W. M.; Ballschmiter, K. (2012). From coal to DDT: the history of the development of the pesticide DDT from synthetic dyes till Silent Spring. *Endeavour*. 36(4): 131–142. https://doi. org/10.1016/j.endeavour.2012.10.003
- Lamberth, C.; Jeanmart, S.; Luksch, T.; Plant, A. (2013). Current Challenges and Trends in the Discovery of Agrochemicals. Science. 341(6147): 742–746. https://doi.org/10.1126/science.1237227
- Liu, N. (2012). Pyrethroid resistance in insects: genes, mechanisms, and regulation. In: Perveen, F.(ed). *Insecticides: Advances in Integrated Pest Management*. pp. 457–468. 1st edition. Croatia: Intechopen. 726p. 10.5772/28737
- Love, R. R.; Sikder, J. R.; Vivero, R. J.; Matute, D. R.; Schrider, D. R. (2023). Strong Positive Selection in *Aedes aegypti* and the Rapid Evolution of Insecticide Resistance. *Molecular Biology and Evolution*. 40(4). https://doi.org/10.1093/molbev/msad072
- Lozano Cruz, B. (1967). *Resistencia de Heliothis virescens Fabricius al D.D.T. en la zona algodonera del Tolima sur*. Ibagué: Universidad del Tolima. 65p.
- Madgwick, P. G.; Wubs, M.; Kanitz, R. (2023). Optimization of long-lasting insecticidal bed nets for resistance management: a modelling study and user-friendly app. *Malaria Journal*. 22(1): 290. https://doi.org/10.1186/s12936-023-04724-x
- Maienfisch, P.; Stevenson, T. M. (2015). Modern Agribusiness Markets, Companies, Benefits and Challenges. In: Maienfisch, P.; Stevenson, T. *Discovery and Synthesis of Crop Protection Products*. pp 1-13. Washington, DC: American Chemical Society. 467 p. https://doi.org/10.1021/ bk-2015-1204.ch001
- Monnerat, R.; Martins, E.; Queiroz, P.; Ordúz, S.; Jaramillo, G.; Benintende, G.; Cozzi, J.; Real, M. D.;
  Martinez-Ramirez, A.; Rausell, C.; Cerón, J.; Ibarra, J. E.; Del Rincon-Castro, M. C.; Espinoza, A. M.; Meza-Basso, L.; Cabrera, L.; Sánchez, J.; Soberon, M.; Bravo, A. (2006). Genetic Variability of Spodoptera frugiperda Smith (Lepidoptera: Noctuidae) Populations from Latin America Is Associated with Variations in Susceptibility to Bacillus thuringiensis Cry Toxins. *Applied and Environmental Microbiology*. 72(11): 7029–7035. https://doi.org/10.1128/AEM.01454-06
- Mota-Sanchez, D.; Wise, J. C. (2024). The arthropod pesticide resistance database. https://www.pesticideresistance.org/search.php
- Murillo, L.; Mosquera, F. (1984). Evaluación de resistencia a tres acaricidas utilizados en el control de Tetranychus cinnabarinus (Boisduval) en cultivos de clavel para exportación. *Revista Colombiana de Entomología*. 10(1): 51–56. https://doi.org/10.25100/socolen.v10il-2.10275
- Navarro, L.; Gongora, C.; Benavides, P. (2010). Single nucleotide polymorphism detection at the *Hypothenemus hampei* Rdl gene by allele-specific PCR amplification with Tm-shift primers. *Pesticide Biochemistry and Physiology*. 97(3): 204–208. https://doi.org/10.1016/j. pestbp.2010.01.006
- Orjuela, L. I.; Morales, J. A.; Ahumada, M. L.; Rios, J. F.; González, J. J.; Yañez, J.; Rosales, A.;



Cabarcas, D. M.; Venegas, J.; Yasnot, M. F.; Quiñones, M. L. (2018). Insecticide Resistance and Its Intensity in populations of malaria vectors in Colombia. *BioMed Research International*. 2018(1): 1–12. https://doi.org/10.1155/2018/9163543

- Ortega, D. S.; Bacca, T.; Nascimento Silva, A. P.; Canal, N. A.; Haddi, K. (2021). Control failure and insecticides resistance in populations of *Rhyzopertha dominica* (Coleoptera: Bostrichidae) from Colombia. *Journal of Stored Products Research*. 92: 101802. https://doi.org/10.1016/j. jspr.2021.101802
- Perry, T.; Batterham, P.; Daborn, P. J. (2011). The biology of insecticidal activity and resistance. *Insect Biochemistry and Molecular Biology*. 41(7): 411–422. https://doi.org/10.1016/j. ibmb.2011.03.003
- Perry, T.; Somers, J.; Yang, Y. T.; Batterham, P. (2015). Expression of insect α6-like nicotinic acetylcholine receptors in *Drosophila melanogaster* highlights a high level of conservation of the receptor: spinosyn interaction. *Insect Biochemistry and Molecular Biology*. 64: 106–115. https://doi.org/10.1016/j.ibmb.2015.01.017
- Ranganathan, M.; Narayanan, M.; Kumarasamy, S. (2022). Importance of Metabolic Enzymes and Their Role in Insecticide Resistance. In: Surajit, M.; Ramkumar, G.; Karthi, S.; Fengliang, J. (eds). New and Future Development in Biopesticide Research: Biotechnological Exploration. pp. 243–260. Singapore: Springer Nature Singapore. https://doi.org/https://doi.org/10.1007/978-981-16-3989-0\_10
- Rendon, F.; Cardona, C. (1976). Aumenta la resistencia del Heliothis a los insecticidas. In: Alvarez-Alcaraz, G. (Ed). Bases técnicas para el cultivo del algodón en Colombia. pp. 384-542. Cuarta edición. Bogotá, D.C. (Colombia): Federación Nacional de Algodoneros. 714 p. https://repository. agrosavia.co
- Rendon, F.; Revelo, R.; Cardona, C. (1977). Resistencia de *Heliothis virescens* a metil paratión y toxametil en nueve zonas algodoneras del país. IV Congreso de La Sociedad Colombiana de Entomología. https://www.socolen.org.co/\_files/ugd/040ab7\_2ab74632cfe243ae9fec7a2c6cbf2ccb.pdf
- Rendon, F.; Revelo, R.; Cardona, C. (1978). La resistencia de plagas del algodonero a los insecticidas de Colombia y sus implicaciones actuales. Manejo de Plaguicidas y Protección Del Ambiente. Seminario sobre manejo de plaguicidas y protección del medio ambiente. https://repository. agrosavia.co/handle/20.500.12324/20798
- Rendón, F.; Siabatto, A.; Herrera, M. A.; Alvarez, A, G. (1990). Results of a pilot programme to monitor the resistance of *Heliothis virescens* (Fabricius) to methyl parathion and pyrethroids in the cotton region of Espinal, 1988 harvest. *Revista Colombiana de Entomologia*. 1(20): 16–22.
- Ríos-Díez, J. D.; Saldamando-Benjumea, C. I. (2011). Susceptibility of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) strains from central Colombia to two insecticides, methomyl and lambda-cyhalothrin: A Study of the genetic basis of resistance. *Journal of Economic Entomology*. 104(5): 1698–1705. https://doi.org/10.1603/EC11079
- Rix, R. R.; Guedes, R. N. C.; Christopher Cutler, G. (2022). Hormesis dose–response contaminantinduced hormesis in animals. *Current Opinion in Toxicology*. 30: 100336. https://doi. org/10.1016/j.cotox.2022.02.009
- Rodriguez-Chalarca, J.; Valencia, S. J.; Rivas-Cano, A.; Santos-González, F.; Romero, D. P. (2024). Impact of Bt corn expressing *Bacillus thuringiensis* Berliner insecticidal proteins on the growth and survival of *Spodoptera frugiperda* larvae in Colombia. Frontiers in Insect Science. 4(1): 1-15. https://doi.org/10.3389/finsc.2024.1268092
- Rodríguez, I.; Cardona, C. (2001). *Problemática de Trialeurodes vaporariorum y Bemisia tabaci* (Homoptera: Aleyrodidae) como plagas de cultivos semestrales en el Valle del Cauca. *Revista Colombiana de Entomología*. 27(1): 21–26. https://doi.org/10.25100/socolen.v27i1.9659
- Rodríguez, I.; Morales, H.; Bueno, J. M.; Cardona, C. (2005). El biotipo B de *Bemisia tabaci* (Homoptera: Aleyrodidae) adquiere mayor importancia en el Valle del Cauca. *Revista Colombiana de Entomología*. 31(1): 21–28.
- Rodríguez, I.; Morales, H.; Cardona, C. (2003). Líneas base, dosis diagnóstica y medición periódica de resistencia a insecticidas en poblaciones de adultos e inmaduros de *Trialeurodes vaporariorum* (Homoptera: Aleyrodidae) en el Valle del Cauca, Colombia. *Revista Colombiana de Entomología*. 29(1): 29–33.
- Rodríguez, I. V.; Bueno, J. M.; Cardona, C.; Morales, H. (2012). *Biotipo B de Bemisia tabaci* (Hemiptera: Aleyrodidae): plaga de pimentón en el Valle del Cauca, Colombia. *Revista*



Colombiana de Entomología. 38(1): 14–22.

- Sajad, K.; Muhammad, N.; Muhammad, R.; Wajid, K.; Muhammad, F.; Shah, A. S.; Fazl, I. S.; Fawad, A.; Khaliq, U. R.; Ahsan, K.; Shahid, A.; Muhammad, M. (2020). Mechanism of Insecticide Resistance in Insects/Pests. *Polish Journal of Environmental Studies*. 29(3): 1–8. https://doi.org/10.15244/pjoes/108513
- Sakuma, M. (1998). Probit analysis of preference data. *Applied Entomology and Zoology*. 33(3): 339–347. https://doi.org/10.1303/aez.33.339
- Sánchez-Bayo, F. (2019). Current-use Pesticides. In: Sanchez-Hernandez., J.(eds). *Bioremediation of Agricultural Soils*. pp. 3–29. 1st Edition. Boca Ratón: CRC Press. 296p. https://doi. org/10.1201/9781315205137-1
- SIB. (2024). Red nacional de datos abiertos sobre biodiversidad. Database. https://biodiversidad. co/data/
- Sparks, T. C.; Crossthwaite, A. J.; Nauen, R., Banba, S.; Cordova, D.; Earley, F.; Ebbinghaus-Kintscher, U.; Fujioka, S.; Hirao, A.; Karmon, D.; Kennedy, R.; Nakao, T.; Popham, H. J. R.; Salgado, V.; Watson, G. B.; Wedel, B. J.; Wessels, F. J. (2020). Insecticides, biologics and nematicides: Updates to IRAC's mode of action classification a tool for resistance management. *Pesticide Biochemistry and Physiology*. 167: 104587. https://doi.org/10.1016/j.pestbp.2020.104587
- Sparks, T. C.; Lorsbach, B. A. (2017). Agrochemical Discovery Building the Next Generation of Insect Control Agents. In: Goross, A. D.; Ozoe, Y.; Coats J. R.(eds). Advances in Agrochemicals: Ion Channels and G Protein-Coupled Receptors (GPCRs) as Targets for Pest Control. pp. 1–17. Washington: American Chemical Society. 141p. https://doi.org/10.1021/bk-2017-1264.ch001
- Sparks, T. C.; Nauen, R. (2015). IRAC: Mode of action classification and insecticide resistance management. *Pesticide Biochemistry and Physiology*. 121: 122-128. https://doi.org/10.1016/j. pestbp.2014.11.014
- Tyler, P. S.; Evans, N. (1981). A tentative method for detecting resistance to gamma-HCH in three bruchid beetles. *Journal of Stored Products Research*. 17(3): 131–135. https://doi. org/10.1016/0022-474X(81)90012-6
- Valencia-Cataño, S. J.; Rodríguez-Chalarca, J.; Blanco, C. A. (2016). Effect of Genetically-Modified Cotton Cultivars on Demographic Parameters of *Spodoptera frugiperda* in Colombia. *Southwestern Entomologist.* 41(4): 963–970. https://doi.org/10.3958/059.041.0426
- Valencia, E.; Plata, C.; Corredor, C.: Cardona, C. (1993). Resistencia enzimática a insecticidas en larvas de *Heliothis virescens* (Fabricius) (Lepidoptera: Noctuidae). *Revista Colombiana de Entomología*. 19(4): 131–138. https://doi.org/10.25100/socolen.v19i4.10069
- Wang, X.; Wang, R. (2024). Insecticide resistance: Monitoring, mechanism and management. *Journal of Applied Entomology*. 148(7): 735–737. https://doi.org/10.1111/jen.13322
- Wolfenbarger, D. A.; Lukefahr, M. J.; Graham, H. M. (1973). LD50 Values of Methyl Parathion and Endrin to Tobacco Budworms and Bollworms Collected in the Americas and Hypothesis on the Spread of Resistance in These Lepidopterans to These Insecticides. *Journal of Economic Entomology*. 66(1): 211–216. https://doi.org/10.1093/jee/66.1.211
- Zalucki, M.; Furlong, M. (2017). Behavior as a mechanism of insecticide resistance: evaluation of the evidence. *Current Opinion in Insect Science*. 21: 19–25. https://doi.org/10.1016/j. cois.2017.05.006
- Zenner de Polanía, I. (1996). Riesgo del uso de insecticidas en agroecosistemas nuevos, con énfasis en resistencia de insectos a insecticidas. https://agris.fao.org/search/en/providers/122610/records/64775f5ba3fd11e4303b8349
- Zenner, I.; Álvarez, J. A.; Mejía, R.; Bayona, M. A. (2005). Influencia de la toxina Cry1Ac del *Bacillus thuringiensis* sobre el desarrollo del cogollero del maíz *Spodoptera frugiperda* (JE Smith). *Revista Actualidad & Divulgación Científica*. 8(2): 129–139.

